

11/28/00

JCS941 U.S. PTO
09/17/2000

Serial Number No. 20350

TOWNSEND and TOWNSEND and CREW LLP
333 California Center, 8th Floor
San Francisco, California 94111-3834
(415) 576-0200ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
Washington, D.C. 20231

Sir:

Transmitted herewith for filing under 37 CFR 1.53(b) is the

- ☐ patent application of
☒ continuation patent application of
☐ divisional patent application of
☐ continuation-in-part patent application of

Inventor(s)/Applicant Identifier: Stephen P.A. Fodor, et al.

For: VERY LARGE SCALE IMMOBILIZED POLYMER SYNTHESIS

- ☒ Please amend this application by adding the following before the first sentence: "This application is a continuation of 09/245,212, filed February 5, 1999, which is a continuation of 09/063,933, filed April 21, 1998, which is a continuation of 08/466,632, filed June 6, 1995, which is a rule 60 division of 08/390,272, filed February 16, 1995, which is a file wrapper continuation of 07/624,120, filed December 6, 1990, now abandoned, which is a continuation-in-part of Application No. 07/492,462, filed March 7, 1990, now U.S. Patent No. 5,143,854, which is a continuation-in-part of 07/362,901, filed June 7, 1989, now abandoned, the disclosure of which is incorporated by reference."

Enclosed are:

- ☒ Return Postcard;
☒ 59 page(s) of specification;
☒ 9 page(s) of claims;
☒ 1 page of Abstract;
☒ 18 sheet(s) of ☐ formal ☒ informal drawing(s);
☒ A ☒ signed ☐ unsigned Declaration & Power of Attorney;
☒ Request to Reference Previously Filed Identical Computer readable Copy According to 37 CFR 1.821(e); and
☒ Communication Under 37 CFR §§ 1.821-1.825 and Preliminary Amendment.

DO NOT CHARGE THE FILING FEE

Please charge Deposit Account No. 20-1430 as follows:

☒ Any additional fees associated with this paper or during the pendency of this application.

- ☐ A check for \$_____ is enclosed.
☒ extra copies of this sheet are enclosed.

Telephone:
(650) 326-2400Facsimile:
(650) 326-2422

Attorney Docket No. 18547-036720US

"Express Mail" Label No. EL 265821898US

Date of Deposit: November 28, 2000

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above, addressed to:

Assistant Commissioner for Patents
Washington, D.C. 20231By: J. LiebeschuetzRespectfully submitted,
TOWNSEND and TOWNSEND and CREW LLP

J. Liebeschuetz
 Joe Liebeschuetz
 Reg No.: 37,505
 Attorneys for Applicant

JCS941 U.S. PTO
09/17/2000

11/23/00


"Express Mail" Label No. EL 265821898US
Date of Deposit November 28, 2000

PATENT
Attorney Docket No.: 018547-036720US
5

I hereby certify that this is being deposited with the United States Postal
Service "Express Mail Post Office to Address" service under 37 CFR 1.10
on the date indicated above and is addressed to:

Assistant Commissioner for Patents
Washington, D.C. 20231

By: _____



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Stephen P.A. Fodor, et al.

Application No.: Unassigned

Filed: November 28, 2000

For: VERY LARGE SCALE
IMMOBILIZED POLYMER
SYNTHESIS

Examiner: Unassigned

Art Unit: Unassigned

COMMUNICATION UNDER

37 C.F.R. §§ 1.821-1.825

AND

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

Sir:

Applicant submits herewith the required paper copy of the Sequence Listing.
Applicant has concurrently filed Request To Reference Previously Filed Identical Computer
Readable Copy According To 37 CFR 1.821(e); and, therefore, Applicant has not submitted
computer readable copy of the sequence listing herewith.

Please amend the specification in adherence with 37 C.F.R. §§ 1.821-1.825 as
follows:

IN THE SPECIFICATION:


Please insert the accompanying paper copy of the Sequence Listing, page
numbers 1 through 22, at the end of the application.

REMARKS

Applicant requests entry of this amendment in adherence with 37 C.F.R. §§1.821 to 1.825. This amendment is accompanied by a paper copy containing the above named sequences, SEQ ID NOS:1-21. The paper copy of the sequence information has been printed from the floppy disk as filed in Application No. 08/466,632. The information contained in the computer readable disk file in application no. 08/466,632 was prepared through the use of the software program "PatentIn" and is identical to that of the paper copy filed herewith. This amendment contains no new matter.

If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 650-326-2400.

Respectfully submitted,



Joe Liebeschuetz
Reg. No. 37,505

TOWNSEND and TOWNSEND and CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, California 94111-3834
Tel: (650) 326-2400
Fax: (650) 326-2422
JOL:llh

PATENT APPLICATION

VERY LARGE SCALE IMMOBILIZED POLYMER SYNTHESIS

Inventor(s):

STEPHEN P.A. FODOR, a citizen of the United States of America, residing in Palo Alto, California;

LUBERT STRYER, a citizen of the United States of America, residing in Stanford, California;

JAMES L. WINKLER, a citizen of the United States of America, residing in _____;

CHRISTOPHER P. HOLMES, a citizen of the United States of America, residing in Sunnyvale, California; and

DENNIS W. SOLAS, a citizen of the United States of America, residing in San Francisco, California.

Assignee:

AFFYMAX TECHNOLOGIES, N.V., a Netherlands Antilles corporation

VERY LARGE SCALE IMMOBILIZED POLYMER SYNTHESIS

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation-in-part
of pending application Serial No. 492,462, filed
March 7, 1990, which is a continuation-in-part of pending
application Serial No. 362,901, filed June 7, 1989,
assigned to the assignee of the present application, and
10 incorporated herein by reference for all purposes. This
application is related to applications with Serial Nos.
_____ and _____, (Attorney Docket
Nos. 11509-26 and 11509-30) filed on the same day as the
present application, and also incorporated herein by
reference for all purposes.

15

COPYRIGHT NOTICE

A portion of the disclosure of this patent
document contains material which is subject to copyright
protection. The copyright owner has no objection to the
20 facsimile reproduction by anyone of the patent document
or the patent disclosure as it appears in the Patent and
Trademark Office patent file or records, but otherwise
reserves all copyright rights whatsoever.

25

BACKGROUND OF THE INVENTION

The present invention relates to the field
of polymer synthesis. More specifically, the invention
provides a reactor system, a masking strategy,
photoremovable protective groups, data collection and
30 processing techniques, and applications for light
directed synthesis of diverse polymer sequences on
substrates.

35


SUMMARY OF THE INVENTION

Methods, apparatus, and compositions for synthesis and use of techniques for diverse polymer sequences on a substrate are disclosed, as well as applications thereof.

According to one aspect of the invention, an improved reactor system for synthesis of diverse polymer sequences on a substrate is provided. According to this embodiment the invention provides for a reactor for contacting reaction fluids to a substrate; a system for delivering selected reaction fluids to the reactor; a translation stage for moving a mask or substrate from at least a first relative location relative to a second relative location; a light for illuminating the substrate through a mask at selected times; and an appropriately programmed digital computer for selectively directing a flow of fluids from the reactor system, selectively activating the translation stage, and selectively illuminating the substrate so as to form a plurality of diverse polymer sequences on the substrate at predetermined locations.

The invention also provides a technique for selection of linker molecules in VLSIPS. According to this aspect of the invention, the invention provides a method of screening a plurality of linker polymers for use in binding affinity studies. The invention includes the steps of forming a plurality of linker polymers on a substrate in selected regions, the linker polymers formed by the steps of recursively: on a surface of a substrate, irradiating a portion of the selected regions to remove a protective group, and contacting the surface with a monomer; contacting the plurality of linker polymers with a ligand; and contacting the ligand with a labeled receptor.

According to another aspect of the invention, improved photoremovable protective groups are provided. According to this aspect of the invention a compound having the formula:



wherein $n = 0$ or 1 ; Y is selected from the group consisting of an oxygen of the carboxyl group of a natural or unnatural amino acid, an amino group of a natural or unnatural amino acid, or the C-5' oxygen group of a natural or unnatural deoxyribonucleic or ribonucleic acid; R^1 and R^2 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfido, or phosphido group; and R^3 is a alkoxy, alkyl, aryl, hydrogen, or alkenyl group is provided.


The invention also provides improved masking techniques for VLSIPS. According to one aspect of the masking technique, the invention provides an ordered method for forming a plurality of polymer sequences by sequential addition of reagents comprising the step of serially protecting and deprotecting portions of the plurality of polymer sequences for addition of other portions of the polymer sequences using a binary synthesis strategy.

Improved data collection equipment and techniques are also provided. According to one embodiment, the instrumentation provides a system for determining affinity of a receptor to a ligand comprising: means for applying light to a surface of a substrate, the substrate comprising a plurality of ligands at predetermined locations, the means for

applying directing light providing simultaneous illumination at a plurality of the predetermined locations; and an array of detectors for detecting light fluoresced at the plurality of predetermined locations.

5 The invention further provides for improved data analysis techniques including the steps of exposing fluorescently labelled receptors to a substrate, the substrate comprising a plurality of ligands in regions at known locations; at a plurality of data collection points
10 within each of the regions, determining an amount of light fluoresced from the data collection points; removing the data collection points deviating from a predetermined statistical distribution; and determining a relative binding affinity of the receptor to remaining
15 data collection points.

Protected amino acid N-carboxy anhydrides for use in polymer synthesis are also disclosed. According to this aspect of the invention provides a compound having the formula:



where R is a side chain of a natural or unnatural amino acid and X is a photoremovable protecting group.

A further understanding of the nature and
30 advantages of the inventions herein may be realized by reference to the remaining portions of the specification and the attached drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 schematically illustrates light-directed spatially-addressable parallel chemical synthesis;

5 Fig. 2 schematically illustrates one example of light-directed peptide synthesis;

Fig. 3 is a three-dimensional representation of a portion of the checkerboard array of YGGFL and PGGFL;

10 Fig. 4 schematically illustrates the software sequences for the automated system for synthesizing diverse polymer sequences;

Fig. 5a and 5b illustrate operation of a program for polymer sythesis;

Fig. 6 is a schematic illustration of a "pure" binary masking strategy;

15 Fig. 7 is a schematic illustration of a gray code binary masking strategy;

Fig. 8 is a schematic illustration of a modified gray code binary masking strategy;

20 Fig. 9a schematically illustrates a masking scheme for a four step synthesis;

Fig. 9b schematically illustrates synthesis of all 400 peptide dimers;

Fig. 10 is a coordinate map for the ten-step binary synthesis;

25 Fig. 11 schematically illustrates a data collection system;

Fig. 12 is a block diagram illustrating the architecture of the data collection system;

30 Fig. 13 is a flow chart illustrating operation of software for the data collection/analysis system; and

Fig. 14 illustrates a three-dimensional plot of intensity versus position for light directed synthesis of a dinucleotide.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

CONTENTS

	I. Definitions
5	II. General
	A. Deprotection and Addition
	1. Example
	2. Example
	B. Antibody recognition
10	1. Example
	III. Synthesis
	A. Reactor System
	B. Binary Synthesis Strategy
15	1. Example
	2. Example
	3. Example
	4. Example
	5. Example
	6. Example
	C. Linker Selection
20	D. Protecting Groups
	1. Use of Photoremovable Groups
	During Solid-Phase Synthesis
	of Peptides
	2. Use of Photoremovable Groups
	During Solid-Phase Synthesis
	of Oligonucleotides
25	E. Amino Acid N-Carboxy
	Anhydrides Protected with
	a Photoremovable Group
	IV. Data Collection
30	A. Data Collection System
	B. Data Analysis
	V. Other Representative Applications
	A. Oligonucleotide Synthesis
35	1. Example
	VI. Conclusion

I. Definitions

Certain terms used herein are intended to have the following general definitions:

- 5 1. Complementary: Refers to the topological compatibility or matching together of interacting surfaces of a ligand molecule and its receptor. Thus, the receptor and its ligand can be described as complementary, and furthermore, the contact
10 surface characteristics are complementary to each other.
- 15 2. Epitope: The portion of an antigen molecule which is delineated by the area of interaction with the subclass of receptors known as antibodies.
- 20 3. Ligand: A ligand is a molecule that is recognized by a particular receptor. Examples of ligands that can be investigated by this invention include, but are not restricted to, agonists and antagonists for cell membrane receptors, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids, etc.), hormone receptors, peptides, enzymes, enzyme
25 substrates, cofactors, drugs, lectins, sugars, oligonucleotides, nucleic acids, oligosaccharides, proteins, and monoclonal antibodies.
- 30 4. Monomer: A member of the set of small molecules which can be joined together to form a polymer. The set of monomers includes but is not restricted to, for example, the set of common L-amino acids, the set of D-amino acids, the set of synthetic amino acids, the set of nucleotides and the set of
35 pentoses and hexoses. As used herein, monomers refers to any member of a basis set for synthesis of a polymer. For example, dimers of the 20 naturally occurring L-amino acids form a basis set of 400

monomers for synthesis of polypeptides. Different basis sets of monomers may be used at successive steps in the synthesis of a polymer. Furthermore, each of the sets may include protected members which are modified after synthesis.

5. Peptide: A polymer in which the monomers are alpha amino acids and which are joined together through amide bonds and alternatively referred to as a polypeptide. In the context of this specification it should be appreciated that the amino acids may be the L-optical isomer or the D-optical isomer. Peptides are often two or more amino acid monomers long, and often more than 20 amino acid monomers long. Standard abbreviations for amino acids are used (e.g., P for proline). These abbreviations are included in Stryer, Biochemistry, Third Ed., 1988, which is incorporated herein by reference for all purposes.
6. Radiation: Energy which may be selectively applied including energy having a wavelength of between 10^{-14} and 10^4 meters including, for example, electron beam radiation, gamma radiation, x-ray radiation, ultra-violet radiation, visible light, infrared radiation, microwave radiation, and radio waves. "Irradiation" refers to the application of radiation to a surface.
7. Receptor: A molecule that has an affinity for a given ligand. Receptors may be naturally-occurring or manmade molecules. Also, they can be employed in their unaltered state or as aggregates with other species. Receptors may be attached, covalently or noncovalently, to a binding member, either directly or via a specific binding substance. Examples of receptors which can be employed by this invention include, but are not restricted to, antibodies,

cell membrane receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants (such as on viruses, cells or other materials), drugs, polynucleotides, nucleic acids, peptides, cofactors, lectins, sugars, polysaccharides, cells, cellular membranes, and organelles. Receptors are sometimes referred to in the art as anti-ligands. As the term receptors is used herein, no difference in meaning is intended. A "Ligand Receptor Pair" is formed when two macromolecules have combined through molecular recognition to form a complex.

Other examples of receptors which can be investigated by this invention include but are not restricted to:

- a) Microorganism receptors: Determination of ligands which bind to receptors, such as specific transport proteins or enzymes essential to survival of microorganisms, is useful in a new class of antibiotics. Of particular value would be antibiotics against opportunistic fungi, protozoa, and those bacteria resistant to the antibiotics in current use.
- b) Enzymes: For instance, the binding site of enzymes such as the enzymes responsible for cleaving neurotransmitters; determination of ligands which bind to certain receptors to modulate the action of the enzymes which cleave the different neurotransmitters is useful in the development of drugs which can be used in the treatment of disorders of neurotransmission.
- c) Antibodies: For instance, the invention may be useful in investigating the ligand-binding site on the antibody molecule which combines with the epitope of an antigen of interest;

determining a sequence that mimics an antigenic epitope may lead to the development of vaccines of which the immunogen is based on one or more of such sequences or lead to the development of related diagnostic agents or compounds useful in therapeutic treatments such as for autoimmune diseases (e.g., by blocking the binding of the "self" antibodies).

- d) Nucleic Acids: Sequences of nucleic acids may be synthesized to establish DNA or RNA binding sequences.
- e) Catalytic Polypeptides: Polymers, preferably polypeptides, which are capable of promoting a chemical reaction involving the conversion of one or more reactants to one or more products. Such polypeptides generally include a binding site specific for at least one reactant or reaction intermediate and an active functionality proximate to the binding site, which functionality is capable of chemically modifying the bound reactant. Catalytic polypeptides are described in, for example, U.S. application Serial No. 404,920, which is incorporated herein by reference for all purposes.
- f) Hormone receptors: For instance, the receptors for insulin and growth hormone. Determination of the ligands which bind with high affinity to a receptor is useful in the development of, for example, an oral replacement of the daily injections which diabetics must take to relieve the symptoms of diabetes, and in the other case, a replacement for the scarce human growth hormone which can only be obtained from cadavers or by recombinant DNA technology. Other examples are the vasoconstrictive hormone receptors; determination of those ligands which

bind to a receptor may lead to the development of drugs to control blood pressure.

- g) Opiate receptors: Determination of ligands which bind to the opiate receptors in the brain is useful in the development of less-addictive replacements for morphine and related drugs.

8. Substrate: A material having a rigid or semi-rigid surface. In many embodiments, at least one surface of the substrate will be substantially flat, although in some embodiments it may be desirable to physically separate synthesis regions for different polymers with, for example, wells, raised regions, etched trenches, or the like. According to other embodiments, small beads may be provided on the surface which may be released upon completion of the synthesis.

9. Protective Group: A material which is chemically bound to a monomer unit and which may be removed upon selective exposure to an activator such as electromagnetic radiation. Examples of protective groups with utility herein include those comprising nitropiperonyl, pyrenylmethoxy-carbonyl, nitroveratryl, nitrobenzyl, dimethyl dimethoxybenzyl, 5-bromo-7-nitroindoliny, o-hydroxy- α -methyl cinnamoyl, and 2-oxymethylene anthraquinone.

10. Predefined Region: A predefined region is a localized area on a surface which is, was, or is intended to be activated for formation of a polymer. The predefined region may have any convenient shape, e.g., circular, rectangular, elliptical, wedge-shaped, etc. For the sake of brevity herein, "predefined regions" are sometimes referred to simply as "regions."

11. Substantially Pure: A polymer is considered to be "substantially pure" within a predefined region of a substrate when it exhibits characteristics that distinguish it from other predefined regions.

5 Typically, purity will be measured in terms of biological activity or function as a result of uniform sequence. Such characteristics will typically be measured by way of binding with a selected ligand or receptor.

10

12. Activator refers to an energy source adapted to render a group active and which is directed from a source to a predefined location on a substrate. A primary illustration of an activator is light. Other examples of activators include ion beams, electric fields, magnetic fields, electron beams, x-ray, and the like.

15

13. Binary Synthesis Strategy refers to an ordered strategy for parallel synthesis of diverse polymer sequences by sequential addition of reagents which may be represented by a reactant matrix, and a switch matrix, the product of which is a product matrix. A reactant matrix is a $1 \times n$ matrix of the building blocks to be added. The switch matrix is all or a subset of the binary numbers, preferably ordered, between 1 and n arranged in columns. In preferred embodiments, a binary strategy is one in which at least two successive steps illuminate half of a region of interest on the substrate. In most preferred embodiments, binary synthesis refers to a synthesis strategy which also factors a previous addition step. For example, a strategy in which a switch matrix for a masking strategy halves regions that were previously illuminated, illuminating about half of the previously illuminated region and protecting the remaining half (while also protecting

20

25

30

35

about half of previously protected regions and illuminating about half of previously protected regions). It will be recognized that binary rounds may be interspersed with non-binary rounds and that only a portion of a substrate may be subjected to a binary scheme, but will still be considered to be a binary masking scheme within the definition herein. A binary "masking" strategy is a binary synthesis which uses light to remove protective groups from materials for addition of other materials such as amino acids. In preferred embodiments, selected columns of the switch matrix are arranged in order of increasing binary numbers in the columns of the switch matrix.

14. Linker refers to a molecule or group of molecules attached to a substrate and spacing a synthesized polymer from the substrate for exposure/binding to a receptor.

II. General

The present invention provides synthetic strategies and devices for the creation of large scale chemical diversity. Solid-phase chemistry, photolabile protecting groups, and photolithography are brought together to achieve light-directed spatially-addressable parallel chemical synthesis in preferred embodiments.

The invention is described herein for purposes of illustration primarily with regard to the preparation of peptides and nucleotides, but could readily be applied in the preparation of other polymers. Such polymers include, for example, both linear and cyclic polymers of nucleic acids, polysaccharides, phospholipids, and peptides having either α -, β -, or ω -amino acids, heteropolymers in which a known drug is covalently bound to any of the above, polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene

sulfides, polysiloxanes, polyimides, polyacetates, or other polymers which will be apparent upon review of this disclosure. It will be recognized further, that illustrations herein are primarily with reference to C- to N-terminal synthesis, but the invention could readily be applied to N- to C-terminal synthesis without departing from the scope of the invention.

A. Deprotection and Addition

The present invention uses a masked light source or other activator to direct the simultaneous synthesis of many different chemical compounds. Fig. 1 is a flow chart illustrating the process of forming chemical compounds according to one embodiment of the invention. Synthesis occurs on a solid support 2. A pattern of illumination through a mask 4a using a light source 6 determines which regions of the support are activated for chemical coupling. In one preferred embodiment activation is accomplished by using light to remove photolabile protecting groups from selected areas of the substrate.

After deprotection, a first of a set of building blocks (indicated by "A" in Fig. 1), each bearing a photolabile protecting group (indicated by "X") is exposed to the surface of the substrate and it reacts with regions that were addressed by light in the preceding step. The substrate is then illuminated through a second mask 4b, which activates another region for reaction with a second protected building block "B". The pattern of masks used in these illuminations and the sequence of reactants define the ultimate products and their locations, resulting in diverse sequences at predefined locations, as shown with the sequences ACEG and BDFH in the lower portion of Fig. 1. Preferred embodiments of the invention take advantage of combinatorial masking strategies to form a large number of compounds in a small number of chemical steps.

A high degree of miniaturization is possible because the density of compounds is determined largely with regard to spatial addressability of the activator, in one case the diffraction of light. Each compound is physically accessible and its position is precisely known. Hence, the array is spatially-addressable and its interactions with other molecules can be assessed.

In a particular embodiment shown in Fig. 1, the substrate contains amino groups that are blocked with a photolabile protecting group. Amino acid sequences are made accessible for coupling to a receptor by removal of the photoprotective groups.

When a polymer sequence to be synthesized is, for example, a polypeptide, amino groups at the ends of linkers attached to a glass substrate are derivatized with nitroveratryloxycarbonyl (NVOC), a photoremovable protecting group. The linker molecules may be, for example, aryl acetylene, ethylene glycol oligomers containing from 2-10 monomers, diamines, diacids, amino acids, or combinations thereof. Photodeprotection is effected by illumination of the substrate through, for example, a mask wherein the pattern has transparent regions with dimensions of, for example, less than 1 cm^2 , 10^{-1} cm^2 , 10^{-2} cm^2 , 10^{-3} cm^2 , 10^{-4} cm^2 , 10^{-5} cm^2 , 10^{-6} cm^2 , 10^{-7} cm^2 , 10^{-8} cm^2 , or 10^{-10} cm^2 . In a preferred embodiment, the regions are between about $10 \times 10 \text{ } \mu\text{m}$ and $500 \times 500 \text{ } \mu\text{m}$. According to some embodiments, the masks are arranged to produce a checkerboard array of polymers, although any one of a variety of geometric configurations may be utilized.

1. Example

In one example of the invention, free amino groups were fluorescently labelled by treatment of the entire substrate surface with fluorescein isothiocyanate (FITC) after photodeprotection. Glass microscope slides were cleaned, aminated by treatment with

0.1% aminopropyltriethoxysilane in 95% ethanol, and incubated at 110°C for 20 min. The aminated surface of the slide was then exposed to a 30 mM solution of the N-hydroxysuccinimide ester of NVOC-GABA (nitroveratryloxycarbonyl- γ -amino butyric acid) in DMF. The NVOC protecting group was photolytically removed by imaging the 365 nm output from a Hg arc lamp through a chrome on glass 100 μ m checkerboard mask onto the substrate for 20 min at a power density of 12 mW/cm². The exposed surface was then treated with 1 mM FITC in DMF. The substrate surface was scanned in an epi-fluorescence microscope (Zeiss Axioskop 20) using 488 nm excitation from an argon ion laser (Spectra-Physics model 2025). The fluorescence emission above 520 nm was detected by a cooled photomultiplier (Hamamatsu 943-02) operated in a photon counting mode. Fluorescence intensity was translated into a color display with red in the highest intensity and black in the lowest intensity areas. The presence of a high-contrast fluorescent checkerboard pattern of 100x100 μ m elements revealed that free amino groups were generated in specific regions by spatially-localized photodeprotection.

2. Example

Fig. 2 is a flow chart illustrating another example of the invention. Carboxy-activated NVOC-leucine was allowed to react with an aminated substrate. The carboxy activated HOBT ester of leucine and other amino acids used in this synthesis was formed by mixing 0.25 mmol of the NVOC amino protected amino acid with 37 mg HOBT (1-hydroxybenzotriazole), 111 mg BOP (benzotriazolyl-n-oxy-tris (dimethylamino)-phosphoniumhexa-fluorophosphate) and 86 μ l DIEA (diisopropylethylamine) in 2.5 ml DMF. The NVOC protecting group was removed by uniform illumination. Carboxy-activated NVOC-phenylalanine was coupled to the exposed amino groups for 2 hours at room temperature,

and then washed with DMF and methylene chloride. Two unmasked cycles of photodeprotection and coupling with carboxy-activated NVOC-glycine were carried out. The surface was then illuminated through a chrome on glass 50 μ m checkerboard pattern mask. Carboxy-activated Na-tBOC-O-tButyl-L-tyrosine was then added. The entire surface was uniformly illuminated to photolyze the remaining NVOC groups. Finally, carboxy-activated NVOC-L-proline was added, the NVOC group was removed by illumination, and the t-BOC and t-butyl protecting groups were removed with TFA. After removal of the protecting groups, the surface consisted of a 50 μ m checkerboard array of Tyr-Gly-Gly-Phe-Leu (YGGFL) and Pro-Gly-Gly-Phe-Leu (PGGFL).

B. Antibody Recognition

In one preferred embodiment the substrate is used to determine which of a plurality of amino acid sequences is recognized by an antibody of interest.

1. Example

In one example, the array of pentapeptides in the example illustrated in Fig. 2 was probed with a mouse monoclonal antibody directed against β -endorphin. This antibody (called 3E7) is known to bind YGGFL and YGGFM with nanomolar affinity and is discussed in Meo et al., Proc. Natl. Acad. Sci. USA (1983) 80:4084, which is incorporated by reference herein for all purposes. This antibody requires the amino terminal tyrosine for high affinity binding. The array of peptides formed as described in Fig. 2 was incubated with a 2 μ g/ml mouse monoclonal antibody (3E7) known to recognize YGGFL. 3E7 does not bind PGGFL. A second incubation with fluoresceinated goat anti-mouse antibody labeled the regions that bound 3E7. The surface was scanned with an epi-fluorescence microscope. The results showed alternating bright and dark 50 μ m squares indicating that

YGGFL and PGGFL were synthesized in geometric array determined by the mask. A high contrast (>12:1 intensity ratio) fluorescence checkerboard image shows that (a) YGGFL and PGGFL were synthesized in alternate 50 μm squares, (b) YGGFL attached to the surface is accessible for binding to antibody 3E7, and (c) antibody 3E7 does not bind to PGGFL.

A three-dimensional representation of the fluorescence intensity data in a portion of the checkerboard is shown in Fig. 3. This figure shows that the border between synthesis sites is sharp. The height of each spike in this display is linearly proportional to the integrated fluorescence intensity in a 2.5 μm pixel. The transition between PGGFL and YGGFL occurs within two spikes (5 μm). There is little variation in the fluorescence intensity of different YGGFL squares. The mean intensity of sixteen YGGFL synthesis sites was 2.03×10^5 counts and the standard deviation was 9.6×10^3 counts.

III. Synthesis

A. Reactor System

Fig. 4 schematically illustrates a device used to synthesize diverse polymer sequences on a substrate. The device includes an automated peptide synthesizer 401. The automated peptide synthesizer is a device which flows selected reagents through a flow cell 402 under the direction of a computer 404. In a preferred embodiment the synthesizer is an ABI Peptide Synthesizer, model no. 431A. The computer may be selected from a wide variety of computers or discrete logic including for, example, an IBM PC-AT or similar computer linked with appropriate internal control systems in the peptide synthesizer. The PC is provided with signals from the board computer indicative of, for example, the end of a coupling cycle.

Substrate 406 is mounted on the flow cell, forming a cavity between the substrate and the flow cell. Selected reagents flow through this cavity from the peptide synthesizer at selected times, forming an array of peptides on the face of the substrate in the cavity. Mounted above the substrate, and preferably in contact with the substrate is a mask 408. Mask 408 is transparent in selected regions to a selected wavelength of light and is opaque in other regions to the selected wavelength of light. The mask is illuminated with a light source 410 such as a UV light source. In one specific embodiment the light source 410 is a model no. 82420 made by Oriel. The mask is held and translated by an x-y-z translation stage 412 such as an x-y translation stage made by Newport Corp. The computer coordinates action of the peptide synthesizer, x-y translation stage, and light source. Of course, the invention may be used in some embodiments with translation of the substrate instead of the mask.

In operation, the substrate is mounted on the reactor cavity. The slide, with its surface protected by a suitable photo removable protective group, is exposed to light at selected locations by positioning the mask and illuminating the light source for a desired period of time (such as, for example, 1 sec to 60 min in the case of peptide synthesis). A selected peptide or other monomer/polymer is pumped through the reactor cavity by the peptide synthesizer for binding at the selected locations on the substrate. After a selected reaction time (such as about 1 sec to 300 min in the case of peptide reactions) the monomer is washed from the system, the mask is appropriately repositioned or replaced, and the cycle is repeated. In most embodiments of the invention, reactions may be conducted at or near ambient temperature.

Figs 5a and 5b are flow charts of the software used in operation of the reactor system. At step 502 the peptide synthesis software is initialized. At step 504 the system calibrates positioners on the x-y translation stage and begins a main loop. At step 506 the system determines which, if any, of the function keys on the computer have been pressed. If F1 has been pressed, the system prompts the user for input of a desired synthesis process. If the user enters F2, the system allows a user to edit a file for a synthesis process at step 510. If the user enters F3 the system loads a process from a disk at step 512. If the user enters F4 the system saves an entered or edited process to disk at step 514. If the user selects F5 the current process is displayed at step 516 while selection of F6 starts the main portion of the program, i.e., the actual synthesis according to the selected process. If the user selects F7 the system displays the location of the synthesized peptides, while pressing F10 returns the user to the disk operating system.

Fig. 5b illustrates the synthesis step 518 in greater detail. The main loop of the program is started in which the system first moves the mask to a next position at step 526. During the main loop of the program, necessary chemicals flow through the reaction cell under the direction of the on-board computer in the peptide synthesizer. At step 528 the system then waits for an exposure command and, upon receipt of the exposure command exposes the substrate for a desired time at step 530. When an acknowledge of exposure complete is received at step 532 the system determines if the process is complete at step 534 and, if so, waits for additional keyboard input at step 536 and, thereafter, exits the perform synthesis process.

A computer program used for operation of the system described above is included as microfiche Appendix A (Copyright, 1990, Affymax Technologies N.V.,

all rights reserved). The program is written in Turbo C++ (Borland Int'l) and has been implemented in an IBM compatible system. The motor control software is adapted from software produced by Newport Corporation. It will be recognized that a large variety of programming languages could be utilized without departing from the scope of the invention herein. Certain calls are made to a graphics program in "Programmer Guide to PC and PS2 Video Systems" (Wilton, Microsoft Press, 1987), which is incorporated herein by reference for all purposes.

Alignment of the mask is achieved by one of two methods in preferred embodiments. In a first embodiment the system relies upon relative alignment of the various components, which is normally acceptable since x-y-z translation stages are capable of sufficient accuracy for the purposes herein. In alternative embodiments, alignment marks on the substrate are coupled to a CCD device for appropriate alignment.

According to some embodiments, pure reagents are not added at each step, or complete photolysis of the protective groups is not provided at each step. According to these embodiments, multiple products will be formed in each synthesis site. For example, if the monomers A and B are mixed during a synthesis step, A and B will bind to deprotected regions, roughly in proportion to their concentration in solution. Hence, a mixture of compounds will be formed in a synthesis region. A substrate formed with mixtures of compounds in various synthesis regions may be used to perform, for example, an initial screening of a large number of compounds, after which a smaller number of compounds in regions which exhibit high binding affinity are further screened. Similar results may be obtained by only partially photolyzing a region, adding a first monomer, re-photolyzing the same region, and exposing the region to a second monomer.

B. Binary Synthesis Strategy

In a light-directed chemical synthesis, the products formed depend on the pattern and order of masks, and on the order of reactants. To make a set of products there will in general be "n" possible masking schemes. In preferred embodiments of the invention herein a binary synthesis strategy is utilized. The binary synthesis strategy is illustrated herein primarily with regard to a masking strategy, although it will be applicable to other polymer synthesis strategies such as the pin strategy, and the like.

In a binary synthesis strategy, the substrate is irradiated with a first mask, exposed to a first building block, irradiated with a second mask, exposed to a second building block, etc. Each combination of masked irradiation and exposure to a building block is referred to herein as a "cycle."

In a preferred binary masking scheme, the masks for each cycle allow irradiation of half of a region of interest on the substrate and protection of the remaining half of the region of interest. By "half" it is intended herein not to mean exactly one-half the region of interest, but instead a large fraction of the region of interest such as from about 30 to 70 percent of the region of interest. It will be understood that the entire masking scheme need not take a binary form; instead non-binary cycles may be introduced as desired between binary cycles.

In preferred embodiments of the binary masking scheme, a given cycle illuminates only about half of the region which was illuminated in a previous cycle, while protecting the remaining half of the illuminated portion from the previous cycle. Conversely, in such preferred embodiments, a given cycle illuminates half of the region which was protected in the previous cycle and protects half the region which was protected in a previous cycle.

In the synthesis strategy disclosed herein, the longest length (ℓ) of the synthesized polymers is $\ell = n/a$; where n is the number of cycles and a is the number of chemical building blocks (note that a given building block may be repeated). The total number of possible compounds synthesized (k) will be $k = a^\ell$. For example, if $a = 20$ and $\ell = 5$, $n = 100$ and $k = 3.2 \times 10^5$.

The synthesis strategy is most readily illustrated and handled in matrix notation. At each synthesis site, the determination of whether to add a given monomer is a binary process. Therefore, each product element P_j is given by the dot product of two vectors, a chemical reactant vector, e.g., $C = [A, B, C, D]$, and a binary vector σ_j . Inspection of the products in the example below for a four-step synthesis, shows that in one four-step synthesis $\sigma_1 = [1, 0, 1, 0]$, $\sigma_2 = [1, 0, 0, 1]$, $\sigma_3 = [0, 1, 1, 0]$, and $\sigma_4 = [0, 1, 0, 1]$, where a 1 indicates illumination and a 0 indicates protection. Therefore, it becomes possible to build a "switch matrix" S from the column vectors σ_j ($j = 1, k$ where k is the number of products).

	σ_1	σ_2	σ_3	σ_4
S	1	1	0	0
	0	0	1	1
	1	0	1	0
	0	1	0	1

The outcome P of a synthesis is simply $P = CS$, the product of the chemical reactant matrix and the switch matrix.

The switch matrix for an n -cycle synthesis yielding k products has n rows and k columns. An important attribute of S is that each row specifies a mask. A two-dimensional mask m_j for the j th chemical step of a synthesis is obtained directly from the j th row of S by placing the elements s_{j1}, \dots, s_{jk} into, for example, a

square format. The particular arrangement below provides a square format, although linear or other arrangements may be utilized.

$$\begin{array}{rcl}
 5 \quad S & = & \begin{array}{cccc} s_{11} & s_{12} & s_{13} & s_{14} \\ s_{21} & s_{22} & s_{23} & s_{24} \\ s_{31} & s_{32} & s_{33} & s_{34} \\ s_{41} & s_{42} & s_{43} & s_{44} \end{array} & m_j = & \begin{array}{cc} s_{j1} & s_{j2} \\ s_{j3} & s_{j4} \end{array}
 \end{array}$$

10 Of course, compounds formed in a light-activated synthesis can be positioned in any defined geometric array. A square or rectangular matrix is convenient but not required. The rows of the switch matrix may be transformed into any convenient array as long as equivalent transformations are used for each row.

15 For example, the masks in the four-step synthesis below are then denoted by:

$$\begin{array}{rcl}
 20 \quad m_1 & = & \begin{array}{cc} 1 & 1 \\ 0 & 0 \end{array} \quad m_2 = \begin{array}{cc} 0 & 0 \\ 1 & 1 \end{array} \quad m_3 = \begin{array}{cc} 1 & 0 \\ 1 & 0 \end{array} \quad m_4 = \begin{array}{cc} 0 & 1 \\ 0 & 1 \end{array}
 \end{array}$$

where 1 denotes illumination (activation) and 0 denotes no illumination.

25 The matrix representation is used to generate a desired set of products and product maps in preferred embodiments. Each compound is defined by the product of the chemical vector and a particular switch vector. Therefore, for each synthesis address, one simply saves the switch vector, assembles all of them into a switch matrix, and extracts each of the rows to form the masks.

30 In some cases, particular product distributions or a maximal number of products are desired. For example, for $C = [A, B, C, D]$, any switch vector (σ_j) consists of four bits. Sixteen four-bit vectors exist. Hence, a maximum of 16 different products can be made by sequential addition of the reagents $[A, B, C, D]$. These 16 column vectors can be assembled in 16! different ways to

form a switch matrix. The order of the column vectors defines the masking patterns, and therefore, the spatial ordering of products but not their makeup. One ordering of these columns gives the following switch matrix (in which "null" (\emptyset) additions are included in brackets for the sake of completeness, although such null additions are elsewhere ignored herein):

	σ_1															σ_{16}	
10		1	1	1	1	1	1	1	0	0	0	0	0	0	0	0	A
		[0	0	0	0	0	0	0	0	1	1	1	1	1	1	1]	\emptyset
	S =	1	1	1	1	0	0	0	0	1	1	1	1	0	0	0	B
		[0	0	0	0	1	1	1	1	0	0	0	0	1	1	1]	\emptyset
		1	1	0	0	1	1	0	0	1	1	0	0	1	1	0	C
15		[0	0	1	1	0	0	1	1	0	0	1	1	0	0	1]	\emptyset
		1	0	1	0	1	0	1	0	1	0	1	0	1	0	1	D
		[0	1	0	1	0	1	0	1	0	1	0	1	0	1]	\emptyset	

The columns of S according to this aspect of the invention are the binary representations of the numbers 15 to 0. The sixteen products of this binary synthesis are ABCD, ABC, ABD, AB, ACD, AC, AD, A, BCD, BC, BD, B, CD, C, D, and \emptyset (null). Also note that each of the switch vectors from the four-step synthesis masks above (and hence the synthesis products) are present in the four bit binary switch matrix. (See columns 6, 7, 10, and 11)

This synthesis procedure provides an easy way for mapping the completed products. The products in the various locations on the substrate are simply defined by the columns of the switch matrix (the first column indicating, for example, that the product ABCD will be present in the upper left-hand location of the substrate). Furthermore, if only selected desired products are to be made, the mask sequence can be derived by extracting the columns with the desired sequences. For example, to form the product set ABCD, ABD, ACD, AD,

BCD, BD, CD, and D, the masks are formed by use of a switch matrix with only the 1st, 3rd, 5th, 7th, 9th, 11th, 13th, and 15th columns arranged into the switch matrix:

5

$$S = \begin{matrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 \\ & & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ & & & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{matrix}$$

10

To form all of the polymers of length 4, the reactant matrix [ABCDABCDABCDABCD] is used. The switch matrix will be formed from a matrix of the binary numbers from 0 to 2^{16} arranged in columns. The columns having four monomers are then selected and arranged into a switch matrix. Therefore, it is seen that the binary switch matrix in general will provide a representation of all the products which can be made from an n-step synthesis, from which the desired products are then extracted.

20

The rows of the binary switch matrix will, in preferred embodiments, have the property that each masking step illuminates half of the synthesis area. Each masking step also factors the preceding masking step; that is, half of the region that was illuminated in the preceding step is again illuminated, whereas the other half is not. Half of the region that was unilluminated in the preceding step is also illuminated, whereas the other half is not. Thus, masking is recursive. The masks are constructed, as described previously, by extracting the elements of each row and placing them in a square array. For example, the four masks in S for a four-step synthesis are:

30

$$\begin{matrix} m_1 = 1 & 1 & 1 & 1 & m_2 = 1 & 1 & 1 & 1 & m_3 = 1 & 1 & 0 & 0 & m_4 = 1 & 0 & 1 & 0 \\ 35 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 \\ & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 \\ & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 \end{matrix}$$

The recursive factoring of masks allows the products of a light-directed synthesis to be represented by a polynomial. (Some light activated syntheses can only be denoted by irreducible, i.e., prime polynomials.)
 5 For example, the polynomial corresponding to the top synthesis of Fig. 9a (discussed below) is

$$P = (A + B)(C + D)$$

- 10 A reaction polynomial may be expanded as though it were an algebraic expression, provided that the order of joining of reactants X_1 and X_2 is preserved ($X_1X_2 \neq X_2X_1$), i.e., the products are not commutative. The product then is $AC + AD + BC + BD$. The polynomial explicitly
 15 specifies the reactants and implicitly specifies the mask for each step. Each pair of parentheses demarcates a round of synthesis. The chemical reactants of a round (e.g., A and B) react at nonoverlapping sites and hence cannot combine with one other. The synthesis area is
 20 divided equally amongst the elements of a round (e.g., A is directed to one-half of the area and B to the other half). Hence, the masks for a round (e.g., the masks m_A and m_B) are orthogonal and form an orthonormal set. The polynomial notation also signifies that each element in a
 25 round is to be joined to each element of the next round (e.g., A with C, A with D, B with C, and B with D). This is accomplished by having m_C overlap m_A and m_B equally, and likewise for m_D . Because C and D are elements of a round, m_C and m_D are orthogonal to each other and form an
 30 orthonormal set.

The polynomial representation of the binary synthesis described above, in which 16 products are made from 4 reactants, is

$$35 \quad P = (A + \emptyset)(B + \emptyset)(C + \emptyset)(D + \emptyset)$$

and the number of chemical steps n is

$$n = \sum b_i$$

- 5 The number of compounds synthesized when $b = a$ and $z = 0$ in all rounds is $a^{n/a}$, compared with 2^n for a binary synthesis. For $n = 20$ and $a = 5$, 625 compounds (all tetrameres) would be formed, compared with 1.049×10^6 compounds in a binary synthesis with the same number of
10 chemical steps.

It should also be noted that rounds in a polynomial can be nested, as in

$$(A + (B+\emptyset)(C+\emptyset)) (D+\emptyset)$$

- 15 The products are AD, BCD, BD, CD, D, A, BC, B, C, and \emptyset .

- Binary syntheses are attractive for two reasons. First, they generate the maximal number of products (2^n) for a given number of chemical steps (n).
20 For four reactants, 16 compounds are formed in the binary synthesis, whereas only 4 are made when each round has two reactants. A 10-step binary synthesis yields 1,024 compounds, and a 20-step synthesis yields 1,048,576. Second, products formed in a binary synthesis are a
25 complete nested set with lengths ranging from 0 to n . All compounds that can be formed by deleting one or more units from the longest product (the n -mer) are present. Contained within the binary set are the smaller sets that would be formed from the same reactants using any other
30 set of masks (e.g., AC, AD, BC, and BD formed in the synthesis shown in Fig. 6 are present in the set of 16 formed by the binary synthesis). In some cases, however, the experimentally achievable spatial resolution may not suffice to accommodate all the compounds formed.
35 Therefore, practical limitations may require one to select a particular subset of the possible switch vectors for a given synthesis.

1. Example

Fig. 6 illustrates a synthesis with binary masking scheme. The binary masking scheme provides the greatest number of sequences for a given number of cycles. According to this embodiment, a mask m1 allows illumination of half of the substrate. The substrate is then exposed to the building block A, which binds at the illuminated regions.

Thereafter, the mask m2 allows illumination of half of the previously illuminated region, while protecting half of the previously illuminated region. The building block B is then added, which binds at the illuminated regions from m2.

The process continues with masks m3, m4, and m5, resulting in the product array shown in the bottom portion of the figure. The process generates 32 (2 raised to the power of the number of monomers) sequences with 5 (the number of monomers) cycles.

2. Example

Fig. 7 illustrates another preferred binary masking scheme which is referred to herein as the gray code masking scheme. According to this embodiment, the masks m1 to m5 are selected such that a side of any given synthesis region is defined by the edge of only one mask. The site at which the sequence BCDE is formed, for example, has its right edge defined by m5 and its left side formed by mask m4 (and no other mask is aligned on the sides of this site). Accordingly, problems created by misalignment, diffusion of light under the mask and the like will be minimized.

3. Example

Fig. 8 illustrates another binary masking scheme. According to this scheme, referred to herein as a modified gray code masking scheme, the number of masks needed is minimized. For example, the mask m2 could be

the same mask as m_1 and simply translated laterally. Similarly, the mask m_4 could be the same as mask m_3 and simply translated laterally.

5 4. Example

A four-step synthesis is shown in Fig. 9a. The reactants are the ordered set $\{A,B,C,D\}$. In the first cycle, illumination through m_1 activates the upper half of the synthesis area. Building block A is then added to give the distribution 602. Illumination through mask m_2 (which activates the lower half), followed by addition of B yields the next intermediate distribution 604. C is added after illumination through m_3 (which activates the left half) giving the distribution 604, and D after illumination through m_4 (which activates the right half), to yield the final product pattern 608 $\{AC,AD,BC,BD\}$.

5. Example

The above masking strategy for the synthesis may be extended for all 400 dipeptides from the 20 naturally occurring amino acids as shown in Fig. 9b. The synthesis consists of two rounds, with 20 photolysis and chemical coupling cycles per round. In the first cycle of round 1, mask 1 activates 1/20th of the substrate for coupling with the first of 20 amino acids. Nineteen subsequent illumination/coupling cycles in round 1 yield a substrate consisting of 20 rectangular stripes each bearing a distinct member of the 20 amino acids. The masks of round 2 are perpendicular to round 1 masks and therefore a single illumination/coupling cycle in round 2 yields 20 dipeptides. The 20 illumination/coupling cycles of round 2 complete the synthesis of the 400 dipeptides.

35 6. Example

The power of the binary masking strategy can be appreciated by the outcome of a 10-step synthesis that

produced 1,024 peptides. The polynomial expression for this 10-step binary synthesis was:

$$(f+\emptyset)(Y+\emptyset)(G+\emptyset)(A+\emptyset)(G+\emptyset)(T+\emptyset)(F+\emptyset)(L+\emptyset)(S+\emptyset)(F+\emptyset)$$

5

Each peptide occupied a 400x400 μm square. A 32x32 peptide array (1,024 peptides, including the null peptide and 10 peptides of $\ell = 1$, and a limited number of duplicates) was clearly evident in a fluorescence scan following side group deprotection and treatment with the antibody 3E7 and fluorescinated antibody. Each synthesis site was a 400x400 μm square.

The scan showed a range of fluorescence intensities, from a background value of 3,300 counts to 22,400 counts in the brightest square ($x = 20$, $y = 9$). Only 15 compounds exhibited an intensity greater than 12,300 counts. The median value of the array was 4,800 counts.

The identity of each peptide in the array could be determined from its x and y coordinate (each range from 0 to 31) and the map of Fig. 10. The chemical units at positions 2, 5, 6, 9, and 10 are specified by the y coordinate and those at positions 1, 3, 4, 7, 8 by the x coordinate. All but one of the peptides was shorter than 10 residues. For example, the peptide at $x = 12$ and $y = 3$ is YGAGF (positions 1, 6, 8, 9, and 10 are nulls). YGAFLS, the brightest element of the array, is at $x = 20$ and $y = 9$.

It is often desirable to deduce a binding affinity of a given peptide from the measured fluorescence intensity. Conceptually, the simplest case is one in which a single peptide binds to a univalent antibody molecule. The fluorescence scan is carried out after the slide is washed with buffer for a defined time. The order of fluorescence intensities is then a measure primarily of the relative dissociation rates of the antibody-peptide complexes. If the on-rate constants are

the same (e.g., if they are diffusion-controlled), the order of fluorescence intensities will correspond to the order of binding affinities. However, the situation is sometimes more complex because a bivalent primary antibody and a bivalent secondary antibody are used. The density of peptides in a synthesis area corresponded to a mean separation of ~7 nm, which would allow multivalent antibody-peptide interactions. Hence, fluorescence intensities obtained according to the method herein will often be a qualitative indicator of binding affinity.

Another important consideration is the fidelity of synthesis. Deletions are produced by incomplete photodeprotection or incomplete coupling. The coupling yield per cycle in these experiments is typically between 85% and 95%. Implementing the switch matrix by masking is imperfect because of light diffraction, internal reflection, and scattering. Consequently, stowaways (chemical units that should not be on board) arise by unintended illumination of regions that should be dark. A binary synthesis array contains many of the controls needed to assess the fidelity of a synthesis. For example, the fluorescence signal from a synthesis area nominally containing a tetrapeptide ABCD could come from a tripeptide deletion impurity such as ACD. Such an artifact would be ruled out by the finding that the fluorescence intensity of the ACD-site is less than that of the ABCD site.

The fifteen most highly labelled peptides in the array obtained with the synthesis of 1,024 peptides described above, were YGAFLS, YGAFS, YGAFL, YGGFSL, YGAF, YGALS, YGGFS, YGAL, YGAFLF, YGAF, YGAFF, YGGLS, YGGFL, YGAFSF, and YGAFLSF. A striking feature is that all fifteen begin with YG, which agrees with previous work showing that an amino-terminal tyrosine is a key determinant of binding. Residue 3 of this set is either A or G, and residue 4 is either F or L. The exclusion of S and T from these positions is clear cut. The finding

that the preferred sequence is YG (A/G) (F/L) fits nicely with the outcome of a study in which a very large library of peptides on phage generated by recombinant DNA methods was screened for binding to antibody 3E7 (see Cwirla et al., Proc. Natl. Acad. Sci. USA, (1990) 87:6378, incorporated herein by reference). Additional binary syntheses based on leads from peptides on phage experiments show that YGAFMQ, YGAFM, and YGAFQ give stronger fluorescence signals than does YGGFM, the immunogen used to obtain antibody 3E7.

Variations on the above masking strategy will be valuable in certain circumstances. For example, if a "kernel" sequence of interest consists of PQR separated from XYZ and that the aim is to synthesize peptides in which these units are separated by a variable number of different residues. The kernel can be placed in each peptide by using a mask that has 1's everywhere. The polynomial representation of a suitable synthesis is:

$$(P) (Q) (R) (A+\emptyset) (B+\emptyset) (C+\emptyset) (D+\emptyset) (X) (Y) (Z)$$

Sixteen peptides will be formed, ranging in length from the 6-mer PQRXYZ to the 10-mer PQRABCDXYZ.

Several other masking strategies will also find value in selected circumstances. By using a particular mask more than once, two or more reactants will appear in the same set of products. For example, suppose that the mask for an 8-step synthesis is

30	A	11110000
	B	00001111
	C	11001100
	D	00110011
	E	10101010
35	F	01010101
	G	11110000
	H	00001111

The products are ACEG, ACFG, ADEG, ADFG, BCEH, BCFH, BDEH, and BDFH. A and G always appear together because their additions were directed by the same mask, and likewise for B and H.

5

C. Linker Selection

According to preferred embodiments the linker molecules used as an intermediary between the synthesized polymers and the substrate are selected for optimum length and/or type for improved binding interaction with a receptor. According to this aspect of the invention diverse linkers of varying length and/or type are synthesized for subsequent attachment of a ligand. Through variations in the length and type of linker, it becomes possible to optimize the binding interaction between an immobilized ligand and its receptor.

The degree of binding between a ligand (peptide, inhibitor, hapten, drug, etc.) and its receptor (enzyme, antibody, etc.) when one of the partners is immobilized on to a substrate will in some embodiments depend on the accessibility of the receptor in solution to the immobilized ligand. The accessibility in turn will depend on the length and/or type of linker molecule employed to immobilize one of the partners. Preferred embodiments of the invention therefore employ the VLSIPS technology described herein to generate an array of, preferably, inactive or inert linkers of varying length and/or type, using photochemical protecting groups to selectively expose different regions of the substrate and to build upon chemically-active groups.

In the simplest embodiment of this concept, the same unit is attached to the substrate in varying multiples or lengths in known locations on the substrate via VLSIPS techniques to generate an array of polymers of varying length. A single ligand (peptide, drug, hapten, etc.) is attached to each of them, and an assay is performed with the binding site to evaluate the degree of

binding with a receptor that is known to bind to the ligand. In cases where the linker length impacts the ability of the receptor to bind to the ligand, varying levels of binding will be observed. In general, the linker which provides the highest binding will then be used to assay other ligands synthesized in accordance with the techniques herein.

According to other embodiments the binding between a single ligand/receptor pair is evaluated for linkers of diverse monomer sequence. According to these embodiments, the linkers are synthesized in an array in accordance with the techniques herein and have different monomer sequence (and, optionally, different lengths). Thereafter, all of the linker molecules are provided with a ligand known to have at least some binding affinity for a given receptor. The given receptor is then exposed to the ligand and binding affinity is deduced. Linker molecules which provide adequate binding between the ligand and receptor are then utilized in screening studies.

D. Protecting Groups

As discussed above, selectively removable protecting groups allow creation of well defined areas of substrate surface having differing reactivities. Preferably, the protecting groups are selectively removed from the surface by applying a specific activator, such as electromagnetic radiation of a specific wavelength and intensity. More preferably, the specific activator exposes selected areas of surface to remove the protecting groups in the exposed areas.

Protecting groups of the present invention are used in conjunction with solid phase oligomer syntheses, such as peptide syntheses using natural or unnatural amino acids, nucleotide syntheses using deoxyribonucleic and ribonucleic acids, oligosaccharide syntheses, and the like. In addition to protecting the substrate surface

from unwanted reaction, the protecting groups block a reactive end of the monomer to prevent self-polymerization. For instance, attachment of a protecting group to the amino terminus of an activated amino acid, such as an N-hydroxysuccinimide-activated ester of the amino acid, prevents the amino terminus of one monomer from reacting with the activated ester portion of another during peptide synthesis.

Alternatively, the protecting group may be attached to the carboxyl group of an amino acid to prevent reaction at this site. Most protecting groups can be attached to either the amino or the carboxyl group of an amino acid, and the nature of the chemical synthesis will dictate which reactive group will require a protecting group.

Analogously, attachment of a protecting group to the 5'-hydroxyl group of a nucleoside during synthesis using for example, phosphate-triester coupling chemistry, prevents the 5'-hydroxyl of one nucleoside from reacting with the 3'-activated phosphate-triester of another.


Regardless of the specific use, protecting groups are employed to protect a moiety on a molecule from reacting with another reagent. Protecting groups of the present invention have the following characteristics: they prevent selected reagents from modifying the group to which they are attached; they are stable (that is, they remain attached to the molecule) to the synthesis reaction conditions; they are removable under conditions that do not adversely affect the remaining structure; and once removed, do not react appreciably with the surface or surface-bound oligomer. The selection of a suitable protecting group will depend, of course, on the chemical nature of the monomer unit and oligomer, as well as the specific reagents they are to protect against.

In a preferred embodiment, the protecting groups are photoactivatable. The properties and uses of photoreactive protecting compounds have been reviewed. See, McCray *et al.*, Ann. Rev. of Biophys. and Biophys.

Chem. (1989) 18:239-270, which is incorporated herein by reference. Preferably, the photosensitive protecting groups will be removable by radiation in the ultraviolet (UV) or visible portion of the electromagnetic spectrum. More preferably, the protecting groups will be removable by radiation in the near UV or visible portion of the spectrum. In some embodiments, however, activation may be performed by other methods such as localized heating, electron beam lithography, laser pumping, oxidation or reduction with microelectrodes, and the like. Sulfonyl compounds are suitable reactive groups for electron beam lithography. Oxidative or reductive removal is accomplished by exposure of the protecting group to an electric current source, preferably using microelectrodes directed to the predefined regions of the surface which are desired for activation. Other methods may be used in light of this disclosure.

Many, although not all, of the photoremovable protecting groups will be aromatic compounds that absorb near-UV and visible radiation. Suitable photoremovable protecting groups are described in, for example, McCray et al., Patchornik, J. Amer. Chem. Soc. (1970) 92:6333, and Amit et al., J. Org. Chem. (1974) 39:192, which are incorporated herein by reference.


A preferred class of photoremovable protecting groups has the general formula:




where R^1 , R^2 , R^3 , and R^4 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido group, or adjacent substituents

(i.e., R^1 - R^2 , R^2 - R^3 , R^3 - R^4) are substituted oxygen groups that together form a cyclic acetal or ketal; R^5 is a hydrogen atom, a alkoxyl, alkyl, hydrogen, halo, aryl, or alkenyl group, and $n = 0$ or 1.


A preferred protecting group, 6-nitroveratryl (NV), which is used for protecting the carboxyl terminus of an amino acid or the hydroxyl group of a nucleotide, for example, is formed when R^2 and R^3 are each a methoxy group, R^1 , R^4 and R^5 are each a hydrogen atom, and $n = 0$:



A preferred protecting group, 6-nitroveratryloxycarbonyl (NVOC), which is used to protect the amino terminus of an amino acid, for example, is formed when R^2 and R^3 are each a methoxy group, R^1 , R^4 and R^5 are each a hydrogen atom, and $n = 1$:




Another preferred protecting group, 6-nitropiperonyl (NP), which is used for protecting the carboxyl terminus of an amino acid or the hydroxyl group of a nucleotide, for example, is formed when R^2 and R^3 together form a methylene acetal, R^1 , R^4 and R^5 are each a hydrogen atom, and $n = 0$:



5

Another preferred protecting group,
 6-nitropiperonyloxycarbonyl (NPOC), which is used to
 protect the amino terminus of an amino acid, for example,
 is formed when R^2 and R^3 together form a methylene acetal,
 R^1 , R^4 and R^5 are each a hydrogen atom, and $n = 1$:


15



20

A most preferred protecting group,
 methyl-6-nitroveratryl (MeNV), which is used for
 protecting the carboxyl terminus of an amino acid or the
 hydroxyl group of a nucleotide, for example, is formed
 when R^2 and R^3 are each a methoxy group, R^1 and R^4 are
 each a hydrogen atom, R^5 is a methyl group, and $n = 0$:


30



Another most preferred protecting group,
 methyl-6-nitroveratryloxycarbonyl (MeNVOC), which is used
 to protect the amino terminus of an amino acid, for
 example, is formed when R^2 and R^3 are each a methoxy

group, R^1 and R^4 are each a hydrogen atom, R^5 is a methyl group, and $n = 1$:


5



10

Another most preferred protecting group, methyl-6-nitropiperonyl (MeNP), which is used for protecting the carboxyl terminus of an amino acid or the hydroxyl group of a nucleotide, for example, is formed when R^2 and R^3 together form a methylene acetal, R^1 and R^4 are each a hydrogen atom, R^5 is a methyl group, and $n = 0$:

15




20

Another most preferred protecting group, methyl-6-nitropiperonyloxycarbonyl (MeNPOC), which is used to protect the amino terminus of an amino acid, for example, is formed when R^2 and R^3 together form a methylene acetal, R^1 and R^4 are each a hydrogen atom, R^5 is a methyl group, and $n = 1$:


25

30

35



A protected amino acid having a photoactivatable oxycarbonyl protecting group, such as NVOC or NPOC or their corresponding methyl derivatives, MeNVOC or MeNPOC, respectively, on the amino terminus is formed by acylating the amine of the amino acid with an activated oxycarbonyl ester of the protecting group. Examples of activated oxycarbonyl esters of NVOC and MeNVOC have the general formula:




NVOC-X

MeNVOC-X

where X is halogen, mixed anhydride, phenoxy, p-nitrophenoxy, N-hydroxysuccinimide, and the like.

A protected amino acid or nucleotide having a photoactivatable protecting group, such as NV or NP or their corresponding methyl derivatives, MeNV or MeNP, respectively, on the carboxy terminus of the amino acid or 5'-hydroxy terminus of the nucleotide, is formed by acylating the carboxy terminus or 5'-OH with an activated benzyl derivative of the protecting group. Examples of activated benzyl derivatives of MeNV and MeNP have the general formula:



MeNV-X

MeNP-X

where X is halogen, hydroxyl, tosyl, mesyl, trifluoromethyl, diazo, azido, and the like.

Another method for generating protected monomers is to react the benzylic alcohol derivative of the protecting group with an activated ester of the

monomer. For example, to protect the carboxyl terminus of an amino acid, an activated ester of the amino acid is reacted with the alcohol derivative of the protecting group, such as 6-nitroveratrol (NVOH). Examples of activated esters suitable for such uses include


5 halo-formate, mixed anhydride, imidazolyl formate, acyl halide, and also includes formation of the activated ester in situ the use of common reagents such as DCC and the like. See Atherton et al. for other examples of

10 activated esters.

A further method for generating protected monomers is to react the benzylic alcohol derivative of the protecting group with an activated carbon of the monomer. For example, to protect the 5'-hydroxyl group of a nucleic acid, a derivative having a 5'-activated carbon is reacted with the alcohol derivative of the protecting group, such as methyl-6-nitropiperonol (MePyROH). Examples of nucleotides having activating groups attached to the 5'-hydroxyl group have the general

15 formula:


20



25

where Y is a halogen atom, a tosyl, mesyl, trifluoromethyl, azido, or diazo group, and the like.


30 Another class of preferred photochemical protecting groups has the formula:




35

where R^1 , R^2 , and R^3 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfanates, sulfido or phosphido group, R^4 and R^5 independently are a hydrogen atom, an alkoxy, alkyl, halo, aryl, hydrogen, or alkenyl group, and $n = 0$ or 1.


A preferred protecting group, 1-pyrenylmethyloxycarbonyl (PyROC), which is used to protect the amino terminus of an amino acid, for example, is formed when R^1 through R^5 are each a hydrogen atom and $n = 1$:



Another preferred protecting group, 1-pyrenylmethyl (PyR), which is used for protecting the carboxy terminus of an amino acid or the hydroxyl group of a nucleotide, for example, is formed when R^1 through R^5 are each a hydrogen atom and $n = 0$:



An amino acid having a pyrenylmethyloxycarbonyl protecting group on its amino terminus is formed by acylation of the free amine of amino acid with an activated oxycarbonyl ester of the pyrenyl protecting group. Examples of activated oxycarbonyl esters of PyROC have the general formula:




5

where X is halogen, or mixed anhydride, p-nitrophenoxy,
or N-hydroxysuccinimide group, and the like.

A protected amino acid or nucleotide having a
photoactivatable protecting group, such as PyR, on the
carboxy terminus of the amino acid or 5'-hydroxy terminus
of the nucleic acid, respectively, is formed by acylating
the carboxy terminus or 5'-OH with an activated
pyrenylmethyl derivative of the protecting group.
Examples of activated pyrenylmethyl derivatives of PyR
have the general formula:

20



25

where X is a halogen atom, a hydroxyl, diazo, or azido
group, and the like.

Another method of generating protected monomers
is to react the pyrenylmethyl alcohol moiety of the
protecting group with an activated ester of the monomer.
For example, an activated ester of an amino acid can be
reacted with the alcohol derivative of the protecting
group, such as pyrenylmethyl alcohol (PyROH), to form the
protected derivative of the carboxy terminus of the amino
acid. Examples of activated esters include halo-formate,
mixed anhydride, imidazolyl formate, acyl halide, and also

35

includes formation of the activated ester in situ and the use of common reagents such as DCC and the like.

Clearly, many photosensitive protecting groups are suitable for use in the present invention.

5 In preferred embodiments, the substrate is irradiated to remove the photoremovable protecting groups and create regions having free reactive moieties and side products resulting from the protecting group. The removal rate of the protecting groups depends on the
10 wavelength and intensity of the incident radiation, as well as the physical and chemical properties of the protecting group itself. Preferred protecting groups are removed at a faster rate and with a lower intensity of radiation. For example, at a given set of conditions,
15 MeNVOC and MeNPOC are photolytically removed from the N-terminus of a peptide chain faster than their unsubstituted parent compounds, NVOC and NPOC, respectively.

Removal of the protecting group is accomplished
20 by irradiation to liberate the reactive group and degradation products derived from the protecting group. Not wishing to be bound by theory, it is believed that irradiation of an NVOC- and MeNVOC-protected oligomers occurs by the following reaction schemes:

25 NVOC-AA \rightarrow 3,4-dimethoxy-6-nitrosobenzaldehyde + CO₂ + AA
MeNVOC-AA \rightarrow 3,4-dimethoxy-6-nitrosoacetophenone + CO₂ + AA

30 where AA represents the N-terminus of the amino acid oligomer.

Along with the unprotected amino acid, other products are liberated into solution: carbon dioxide and a 2,3-dimethoxy-6-nitrosophenylcarbonyl compound, which
35 can react with nucleophilic portions of the oligomer to form unwanted secondary reactions. In the case of an NVOC-protected amino acid, the degradation product is a nitrosobenzaldehyde, while the degradation product for

the other is a nitrosophenyl ketone. For instance, it is believed that the product aldehyde from NVOC degradation reacts with free amines to form a Schiff base (imine) that affects the remaining polymer synthesis. Preferred photoremovable protecting groups react slowly or reversibly with the oligomer on the support.

Again not wishing to be bound by theory, it is believed that the product ketone from irradiation of a MeNVOC-protected oligomer reacts at a slower rate with nucleophiles on the oligomer than the product aldehyde from irradiation of the same NVOC-protected oligomer. Although not unambiguously determined, it is believed that this difference in reaction rate is due to the difference in general reactivity between aldehyde and ketones towards nucleophiles due to steric and electronic effects.

The photoremovable protecting groups of the present invention are readily removed. For example, the photolysis of N-protected L-phenylalanine in solution and having different photoremovable protecting groups was analyzed, and the results are presented in the following table:

Table
Photolysis of Protected L-Phe-OH

Solvent	<u>t_{1/2} in seconds</u>			
	NBOC	NVOC	MeNVOC	MeNPOC
Dioxane	1288	110	24	19
5mM H ₂ SO ₄ /Dioxane	1575	98	33	22

The half life, $t_{1/2}$, is the time in seconds required to remove 50% of the starting amount of protecting group. NBOC is the 6-nitrobenzyloxycarbonyl group, NVOC is the 6-nitroveratryloxycarbonyl group, MeNVOC is the methyl-6-nitroveratryloxycarbonyl group,

and MeNPOC is the methyl-6-nitropiperonyloxycarbonyl group. The photolysis was carried out in the indicated solvent with 362/364 nm-wavelength irradiation having an intensity of 10 mW/cm², and the concentration of each protected phenylalanine was 0.10 mM.

The table shows that deprotection of NVOC-, MeNVOC-, and MeNPOC-protected phenylalanine proceeded faster than the deprotection of NBOC. Furthermore, it shows that the deprotection of the two derivatives that are substituted on the benzylic carbon, MeNVOC and MeNPOC, were photolyzed at the highest rates in both dioxane and acidified dioxane.

1. Use of Photoremovable Groups During Solid-Phase Synthesis of Peptides

The formation of peptides on a solid-phase support requires the stepwise attachment of an amino acid to a substrate-bound growing chain. In order to prevent unwanted polymerization of the monomeric amino acid under the reaction conditions, protection of the amino terminus of the amino acid is required. After the monomer is coupled to the end of the peptide, the N-terminal protecting group is removed, and another amino acid is coupled to the chain. This cycle of coupling and deprotecting is continued for each amino acid in the peptide sequence. See Merrifield, J. Am. Chem. Soc. (1963) 85:2149, and Atherton et al., "Solid Phase Peptide Synthesis" 1989, IRL Press, London, both incorporated herein by reference for all purposes. As described above, the use of a photoremovable protecting group allows removal of selected portions of the substrate surface, via patterned irradiation, during the deprotection cycle of the solid phase synthesis. This selectively allows spatial control of the synthesis--the next amino acid is coupled only to the irradiated areas.

In one embodiment, the photoremovable protecting groups of the present invention are attached to an activated ester of an amino acid at the amino terminus:



10

where R is the side chain of a natural or unnatural amino acid, X is a photoremovable protecting group, and Y is an activated carboxylic acid derivative. The photoremovable protecting group, X, is preferably NVOC, NPOC, PyROC, MeNVOC, MeNPOC, and the like as discussed above. The activated ester, Y, is preferably a reactive derivative having a high coupling efficiency, such as an acyl halide, mixed anhydride, N-hydroxysuccinimide ester, perfluorophenyl ester, or urethane protected acid, and the like. Other activated esters and reaction conditions are well known (See Atherton et al.).


2. Use of Photoremovable Groups During Solid-Phase Synthesis of Oligonucleotides

The formation of oligonucleotides on a solid-phase support requires the stepwise attachment of a nucleotide to a substrate-bound growing oligomer. In order to prevent unwanted polymerization of the monomeric nucleotide under the reaction conditions, protection of the 5'-hydroxyl group of the nucleotide is required. After the monomer is coupled to the end of the oligomer, the 5'-hydroxyl protecting group is removed, and another nucleotide is coupled to the chain. This cycle of coupling and deprotecting is continued for each nucleotide in the oligomer sequence. See Gait, "Oligonucleotide Synthesis: A Practical Approach" 1984, IRL Press, London, incorporated herein by reference for

all purposes. As described above, the use of a photoremovable protecting group allows removal, via patterned irradiation, of selected portions of the substrate surface during the deprotection cycle of the solid phase synthesis. This selectively allows spatial control of the synthesis--the next nucleotide is coupled only to the irradiated areas.

Oligonucleotide synthesis generally involves coupling an activated phosphorous derivative on the 3'-hydroxyl group of a nucleotide with the 5'-hydroxyl group of an oligomer bound to a solid support. Two major chemical methods exist to perform this coupling: the phosphate-triester and phosphoamidite methods (See Gait). Protecting groups of the present invention are suitable for use in either method.


In a preferred embodiment, a photoremovable protecting group is attached to an activated nucleotide on the 5'-hydroxyl group:




where B is the base attached to the sugar ring; R is a hydrogen atom when the sugar is deoxyribose or R is a hydroxyl group when the sugar is ribose; P represents an activated phosphorous group; and X is a photoremovable protecting group. The photoremovable protecting group, X, is preferably NV, NP, PyR, MeNV, MeNP, and the like as described above. The activated phosphorous group, P, is preferably a reactive derivative having a high coupling efficiency, such as a phosphate-triester, phosphoamidite or the like. Other activated phosphorous derivatives, as well as reaction conditions, are well known (See Gait).

E. Amino Acid N-Carboxy Anhydrides
Protected With a Photoremovable Group

During Merrifield peptide synthesis, an activated ester of one amino acid is coupled with the free amino terminus of a substrate-bound oligomer. Activated esters of amino acids suitable for the solid phase synthesis include halo-formate, mixed anhydride, imidazolyl formate, acyl halide, and also includes formation of the activated ester in situ and the use of common reagents such as DCC and the like (See Atherton et al.). A preferred protected and activated amino acid has the general formula:



where R is the side chain of the amino acid and X is a photoremovable protecting group. This compound is a urethane-protected amino acid having a photoremovable protecting group attach to the amine. A more preferred activated amino acid is formed when the photoremovable protecting group has the general formula:




where R^1 , R^2 , R^3 , and R^4 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido group, or adjacent substituents (i.e., R^1-R^2 , R^2-R^3 , R^3-R^4) are substituted oxygen groups

that together form a cyclic acetal or ketal; and R^5 is a hydrogen atom, an alkoxy, alkyl, hydrogen, halo, aryl, or alkenyl group.

A preferred activated amino acid is formed when the photoremovable protecting group is 6-nitroveratryloxycarbonyl. That is, R^1 and R^4 are each a hydrogen atom, R^2 and R^3 are each a methoxy group, and R^5 is a hydrogen atom. Another preferred activated amino acid is formed when the photoremovable group is 6-nitropiperonyl: R^1 and R^4 are each a hydrogen atom, R^2 and R^3 together form a methylene acetal, and R^5 is a hydrogen atom. Other protecting groups are possible. Another preferred activated ester is formed when the photoremovable group is methyl-6-nitroveratryl or methyl-6-nitropiperonyl.

Another preferred activated amino acid is formed when the photoremovable protecting group has the general formula:



where R^1 , R^2 , and R^3 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfonates, sulfido or phosphido group, and R^4 and R^5 independently are a hydrogen atom, an alkoxy, alkyl, halo, aryl, hydrogen, or alkenyl group. The resulting compound is a urethane-protected amino acid having a pyrenylmethyloxycarbonyl protecting group attached to the amine. A more preferred embodiment is formed when R^1 through R^5 are each a hydrogen atom.

The urethane-protected amino acids having a photoremovable protecting group of the present invention

are prepared by condensation of an N-protected amino acid with an acylating agent such as an acyl halide, anhydride, chloroformate and the like (See Fuller et al., U.S. Patent No. 4,946,942 and Fuller et al., J. Amer. Chem. Soc. (1990) 112:7414-7416, both herein incorporated by reference for all purposes).

Urethane-protected amino acids having photoremovable protecting groups are generally useful as reagents during solid-phase peptide synthesis, and because of the spatially selectivity possible with the photoremovable protecting group, are especially useful for the spatially addressing peptide synthesis. These amino acids are difunctional: the urethane group first serves to activate the carboxy terminus for reaction with the amine bound to the surface and, once the peptide bond is formed, the photoremovable protecting group protects the newly formed amino terminus from further reaction. These amino acids are also highly reactive to nucleophiles, such as deprotected amines on the surface of the solid support, and due to this high reactivity, the solid-phase peptide coupling times are significantly reduced, and yields are typically higher.

IV. Data Collection

A. Data Collection System

Substrates prepared in accordance with the above description are used in one embodiment to determine which of the plurality of sequences thereon bind to a receptor of interest. Fig. 11 illustrates one embodiment of a device used to detect regions of a substrate which contain florescent markers. This device would be used, for example, to detect the presence or absence of a labeled receptor such as an antibody which has bound to a synthesized polymer on a substrate.

Light is directed at the substrate from a light source 1002 such as a laser light source of the type well known to those of skill in the art such as a model no.

2025 made by Spectra Physics. Light from the source is directed at a lens 1004 which is preferably a cylindrical lens of the type well known to those of skill in the art. The resulting output from the lens 1004 is a linear beam rather than a spot of light, resulting in the capability to detect data substantially simultaneously along a linear array of pixels rather than on a pixel-by-pixel basis. It will be understood that which a cylindrical lens is used herein as an illustration of one technique for generating a linear beam of light on a surface, it will be understood that other techniques could also be utilized.

The beam from the cylindrical lens is passed through a dichroic mirror or prism and directed at the surface of the suitably prepared substrate 1008. Substrate 1008 is placed on an x-y translation stage 1009 such as a model no. PM500-8 made by Newport. Light at certain locations on the substrate will be fluoresced and transmitted along the path indicated by dashed lines back through the dichroic mirror, and focused with a suitable lens 1010 such as an f/1.4 camera lens on a linear detector 1012 via a variable f stop focusing lens 1014. Through use of a linear light beam, it becomes possible to generate data over a line of pixels (such as about 1 cm) along the substrate, rather than from individual points on the substrate. In alternative embodiments, light is directed at a 2-dimensional area of the substrate and fluoresced light detected by a 2-dimensional CCD array. Linear detection is preferred because substantially higher power densities are obtained.

Detector 1012 detects the amount of light fluoresced from the substrate as a function of position. According to one embodiment the detector is a linear CCD array of the type commonly known to those of skill in the art. The x-y translation stage, the light source, and the detector 1012 are all operably connected to a

computer 1016 such as a IBM PC-AT or equivalent for control of the device and data collection from the CCD array.

5 In operation, the substrate is appropriately positioned by the translation stage. The light source is then illuminated, and intensity data are gathered with the computer via the detector.

Fig. 12 illustrates the architecture of the data collection system in greater detail. Operation of the system occurs under the direction of the photon counting program 1102 (photon), included herewith as Appendix B. The user inputs the scan dimensions, the number of pixels or data points in a region, and the scan speed to the counting program. Via a GPIB bus 1104 the program (in an IBM PC compatible computer, for example) 15 interfaces with a multichannel scaler 1106 such as a Stanford Research SR 430 and an x-y stage controller 1108 such as a PMS500. The signal from the light from the fluorescing substrate enters a photon counter 1110, 20 providing output to the scaler 1106. Data are output from the scaler indicative of the number of counts in a given region. After scanning a selected area, the stage controller is activated with commands for acceleration and velocity, which in turn drives the scan stage 1112 25 such as a PMS500-A to another region.

Data are collected in an image data file 1114 and processed in a scaling program 1116, also included in Appendix B. A scaled image is output for display on, for example, a VGA display 1118. The image is scaled based 30 on an input of the percentage of pixels to clip and the minimum and maximum pixel levels to be viewed. The system outputs for use the min and max pixel levels in the raw data.

B. Data Analysis

The output from the data collection system is an array of data indicative of fluorescent intensity versus location on the substrate. The data are typically
5 taken over regions substantially smaller than the area in which synthesis of a given polymer has taken place. Merely by way of example, if polymers were synthesized in squares on the substrate having dimensions of 500 microns by 500 microns, the data may be taken over regions having
10 dimensions of 5 microns by 5 microns. In most preferred embodiments, the regions over which fluorescence data are taken across the substrate are less than about 1/2 the area of the regions in which individual polymers are synthesized, preferably less than 1/10 the area in which
15 a single polymer is synthesized, and most preferably less than 1/100 the area in which a single polymer is synthesized. Hence, within any area in which a given polymer has been synthesized, a large number of fluorescence data points are collected.

20 A plot of number of pixels versus intensity for a scan of a cell when it has been exposed to, for example, a labeled antibody will typically take the form of a bell curve, but spurious data are observed, particularly at higher intensities. Since it is
25 desirable to use an average of fluorescent intensity over a given synthesis region in determining relative binding affinity, these spurious data will tend to undesirably skew the data.

Accordingly, in one embodiment of the invention
30 the data are corrected for removal of these spurious data points, and an average of the data points is thereafter utilized in determining relative binding efficiency.

Fig. 13 illustrates one embodiment of a system for removal of spurious data from a set of fluorescence
35 data such as data used in affinity screening studies. A user or the system inputs data relating to the chip location and cell corners at step 1302. From this

information and the image file, the system creates a computer representation of a histogram at step 1304, the histogram (at least in the form of a computer file) plotting number of data pixels versus intensity.

5 For each cell, a main data analysis loop is then performed. For each cell, at step 1306, the system calculates the total intensity or number of pixels for the bandwidth centered around varying intensity levels. For example, as shown in the plot to the right of step 10 1306, the system calculates the number of pixels within the band of width w . The system then "moves" this bandwidth to a higher center intensity, and again calculates the number of pixels in the bandwidth. This process is repeated until the entire range of intensities 15 have been scanned, and at step 1308 the system determines which band has the highest total number of pixels. The data within this bandwidth are used for further analysis. Assuming the bandwidth is selected to be reasonably small, this procedure will have the effect of eliminating 20 spurious data located at the higher intensity levels. The system then repeats at step 1310 if all cells have been evaluated, or repeats for the next cell.

At step 1312 the system then integrates the data within the bandwidth for each of the selected cells, 25 sorts the data at step 1314 using the synthesis procedure file, and displays the data to a user on, for example, a video display or a printer.

V. Representative Applications

30 A. Oligonucleotide Synthesis

The generality of light directed spatially addressable parallel chemical synthesis is demonstrated by application to nucleic acid synthesis.

1. Example

Light activated formation of a thymidine-cytidine dimer was carried out. A three dimensional representation of a fluorescence scan showing a checkboard pattern generated by the light-directed synthesis of a dinucleotide is shown in Fig. 8. 5'-nitroveratryl thymidine was attached to a synthesis substrate through the 3' hydroxyl group. The nitroveratryl protecting groups were removed by illumination through a 500 mm checkerboard mask. The substrate was then treated with phosphoramidite activated 2'-deoxycytidine. In order to follow the reaction fluorometrically, the deoxycytidine had been modified with an Fmoc protected aminohexyl linker attached to the exocyclic amine (5'-O-dimethoxytrityl-4-N-(6-N-fluorenylmethylcarbamoyl-hexylcarboxy)-2'-deoxycytidine). After removal of the Fmoc protecting group with base, the regions which contained the dinucleotide were fluorescently labelled by treatment of the substrate with 1 mM FITC in DMF for one hour.

The three-dimensional representation of the fluorescent intensity data in Fig. 14 clearly reproduces the checkerboard illumination pattern used during photolysis of the substrate. This result demonstrates that oligonucleotides as well as peptides can be synthesized by the light-directed method.

VI. Conclusion

The inventions herein provide a new approach for the simultaneous synthesis of a large number of compounds. The method can be applied whenever one has chemical building blocks that can be coupled in a solid-phase format, and when light can be used to generate a reactive group.

The above description is illustrative and not restrictive. Many variations of the invention will become apparent to those of skill in the art upon review

of this disclosure. Merely by way of example, while the invention is illustrated primarily with regard to peptide and nucleotide synthesis, the invention is not so limited. The scope of the invention should, therefore, be determined not with reference to the above description, but instead should be determined with reference to the appended claims along with their full scope of equivalents.

10

15

20

25

30

35

WHAT IS CLAIMED IS:

1. A reactor system for synthesizing a plurality of polymer sequences on a substrate comprising:

- 5 a) a reactor for contacting reaction fluids to said substrate;
- b) a system for delivering selected reaction fluids to said reactor;
- 10 c) a translation stage for moving a mask or substrate from at least a first relative location relative to a second relative location;
- d) a light for illuminating said substrate through a mask at selected times; and
- 15 e) an appropriately programmed digital computer for selectively directing a flow of fluids from said reactor system, selectively activating said translation stage, and selectively illuminating said substrate so as to form a plurality of diverse polymer sequences on said substrate at predetermined locations.

20

2. The reactor system as recited in claim 1 adapted to provide a plurality of monomers in a reaction fluid to said substrate, said substrate used for an initial screening of polymer sequences.

25

3. An ordered method for forming a plurality of polymer sequences by sequential addition of reagents comprising the step of serially protecting and deprotecting portions of said plurality of polymer sequences for addition of other portions of said polymer sequences using a binary synthesis strategy.

30

35

4. The method as recited in claim 3 wherein said binary synthesis strategy is a binary masking strategy.

5. The method as recited in claim 4 wherein said masking strategy in which said masking strategy provides

at least two consecutive steps in which a mask factors a previous mask by protecting a portion of a previously illuminated portions to light and exposing a portion of a previously protected portions to light.

5

6. The method as recited in claim 4 in which said masking strategy in which at least two successive steps in said masking strategy illuminate about one half of a region of interest on said substrate.

10

7. The method as recited in claim 4 wherein said masking strategy forms a plurality of polymer sequences on a single substrate.

15

8. The method as recited in claim 4 wherein said masks are arranged in a gray code masking scheme, said gray code masking scheme having one edge illumination on each of a plurality of synthesis sites.

20

9. The method as recited in claim 4 wherein said masking scheme results in a minimum number of masking steps for a number of polymers synthesized.

25

10. The method as recited in claim 4 wherein all possible polymers of length 1 are formed with a given basis set of monomers.

30

11. The method as recited in claim 4 wherein said masking strategy is developed in an appropriately programmed digital computer inputting at least a desired basis set, and length of polymers.

35

12. The method as recited in claim 4 wherein all possible polymers of a length less than or equal to 1 are formed with a given basis set of monomers.

13. The method as recited in claim 4 further comprising the step of forming a portion of said polymers with a non-binary masking scheme.

5 14. The method as recited in claim 10 further comprising the step of outputting a masking strategy.

10 15. The method as recited in claim 10 further comprising the step of outputting a map of synthesized polymers on said substrate.

16. The method as recited in claim 15 wherein said map is in the form of Fig. 10.

15 17. A method of screening a plurality of linker polymers for use in binding affinity studies comprising the steps of:

- 20 a) forming a plurality of linker polymers on a substrate in selected regions, said linker polymers formed by the steps of recursively:
- i) on a surface of a substrate, irradiating a portion of said selected regions to remove a protective group; and
 - 25 ii) contacting said surface with a monomer;
 - 25 b) contacting said plurality of linker polymers with a ligand; and
 - 30 c) contacting said ligand with a labeled receptor.

30 18. The method as recited in claim 17 wherein said ligand is a polypeptide.

35 19. The method as recited in claim 17 wherein said receptor is an antibody.

20. The method as recited in claim 17 wherein said monomers added in step ii) are the same in each of said

recursive steps, said selected regions comprising linker molecules of different lengths.

21. The method as recited in claim 17 wherein said
5 labelled receptor is a fluoresceinated receptor.

22. A system for determining affinity of a receptor
to a ligand comprising:

10 a) means for applying light to a surface of a
substrate, said substrate comprising a plurality of
ligands at predetermined locations, said means for
applying directing light providing simultaneous
illumination at a plurality of said predetermined
locations; and

15 b) an array of detectors for detecting light
fluoresced at said plurality of predetermined locations.

23. A system as recited in claim 22 wherein said
means for applying light comprises a point light source
20 and a cylindrical lens for focusing said point light
source along a substantially linear path.

24. A system as recited in claim 22 wherein said
array of detectors comprises a linear array.

25 25. A system as recited in claim 22 wherein said
array of detectors comprises a linear CCD array.

26. In a digital computer, a method of determining
30 the tendency of a receptor to bind to a ligand
comprising:

a) exposing fluorescently labelled receptors to
a substrate, said substrate comprising a plurality of
ligands in regions at known locations;


35 b) at a plurality of data collection points
within each of said regions, determining an amount of
light fluoresced from said data collection points;

c) removing said data collection points deviating from a preset amount from a predetermined statistical distribution; and

d) determining a relative binding affinity of said receptor to remaining data collection points.

27. The method as recited in claim 26 wherein said predetermined statistical distribution is a normal distribution.

28. A compound having the formula:



wherein $n = 0$ or 1 ; Y is selected from the group consisting of an oxygen of the carboxyl group of a natural or unnatural amino acid, an amino group of a natural or unnatural amino acid, or the C-5' oxygen group of a natural or unnatural deoxyribonucleic or ribonucleic acid; R^1 and R^2 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfido, or phosphido group; and R^3 is a alkoxy, alkyl, aryl, hydrogen, or alkenyl group.

29. The compound of claim 28 wherein Y is the C-5' oxygen group of a natural or unnatural deoxyribonucleic or ribonucleic acid.

30. The compound of claim 29 wherein $n = 0$.

31. The compound of claim 29 wherein R^1 and R^2 are each a hydrogen atom.

32. The compound of claim 31 wherein R^3 is a hydrogen atom.

33. The compound of claim 31 wherein R^3 is a methyl group.


34. The compound of claim 28 wherein Y is an oxygen of the carboxyl group of an amino acid and $n = 0$.

35. The compound of claim 34 wherein R^1 and R^2 are each a hydrogen atom.

36. The compound of claim 35 wherein R^3 is a hydrogen atom.

37. The compound of claim 35 wherein R^3 is a methyl group.

38. A compound having the formula:



wherein $n = 0$ or 1 ; Y is selected from the group consisting of an amino group of a natural or unnatural amino acid or the C-5' oxygen group of a natural or unnatural deoxyribonucleic and ribonucleic acid; R^1 , R^2 , and R^3 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfido or phosphido group; R^4 and R^5 independently are a


alkoxy, alkyl, hydrogen, halo, aryl, hydrogen, or alkenyl group.

39. The compound of claim 38 wherein R^1 through R^3 are each a hydrogen atom.

40. The compound of claim 39 wherein R^4 and R^5 are each a hydrogen atom.

41. The compound of claim 39 wherein R^4 and R^5 are each a methyl group.

42. A compound having the formula:




wherein $n = 0$ or 1 ; Y is a C-5' oxygen group of a natural or unnatural deoxyribonucleic and ribonucleic acid; R^1 through R^4 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfido, or phosphido group; and R^5 is a alkoxy, alkyl, aryl, or alkenyl group.

43. The compound of claim 42 wherein R^2 and R^3 are each a methoxy group.

44. The compound of claim 43 wherein R^1 and R^4 are each a hydrogen atom.

45. The compound of claim 44 wherein R^5 is a methyl group.

46. A compound having the formula:




wherein $n = 0$ or 1 ; Y is an atom to be protected; R^1 and R^2 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido, sulfido, or phosphido group; and R^3 is a alkoxy, alkyl, aryl, or alkenyl group.

47. The compound of claim 46 wherein Y is selected from the group consisting of an oxygen of the carboxyl group of a natural or unnatural amino acid, or the C-5' oxygen group of a natural or unnatural deoxyribonucleic or ribonucleic acid, or the amino group of a natural or unnatural amino acid.

48. The compound of claim 47 wherein R^1 and R^2 are hydrogen.


49. The compound of claim 48 wherein R^3 is a methyl group.

50. A compound having the formula:



where R is a side chain of a natural or unnatural amino acid and X is a photoremovable protecting group.

51. The compound of claim 50 wherein X has the following formula:



10

where R^1 , R^2 , R^3 , and R^4 independently are a hydrogen atom, a lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido group, or adjacent substituents are substituted oxygen groups that together form a cyclic acetal or ketal; and R^5 is a hydrogen atom, a alkoxy, alkyl, hydrogen, halo, aryl, or alkenyl group.

20

52. The compound of claim 51 wherein R^1 and R^4 are each a hydrogen atom, and R^2 and R^3 are each a methoxy group.

25

53. The compound of claim 52 wherein R^5 is a methyl group.

30

54. The compound of claim 51 wherein R^2 and R^3 are substituted oxygen groups that together form a cyclic acetal.

55. The compound of claim 54 wherein R^1 and R^4 are each a hydrogen atom.

35

56. The compound of claim 55 wherein R^5 is a methyl group.

VERY LARGE SCALE IMMOBILIZED POLYMER SYNTHESIS

5 ABSTRACT OF THE DISCLOSURE

A synthetic strategy for the creation of large
scale chemical diversity. Solid-phase chemistry,
photolabile protecting groups, and photolithography are
used to achieve light-directed spatially-addressable
10 parallel chemical synthesis. Binary masking techniques
are utilized in one embodiment. A reactor system,
photoremovable protective groups, and improved data
collection and handling techniques are also disclosed. A
technique for screening linker molecules is also
15 provided.

20 WP50/11509/028.APP

25

30

35




FIG. 1.

- a) $h\nu$; x-L
- b) $h\nu$; x-F
- c) $h\nu$; x-G
- d) $h\nu$; x-G

e) \overline{Y}

f) 
p

FIG. 2.

1000




FIG. 5A.




FIG. 5B.

074403-1-000000




FIG. 6A.

FIG. 6A.

FIG. 6B.

FIG. 6.

SECRET 100-440




FIG. 6B.

FIG. 7A





FIG. 7A.

FIG. 7A.
FIG. 7B.

FIG. 7.



09724001-1 100000




FIG. 8A.

FIG. 8A.
FIG. 8B.

FIG. 8






FIG. 8B.



(x, y) [illegible]

FIG. 10.





FIG. 11.






FIG. 14.




FIG. 14.

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

IMPROVED POLYMER SYNTHESIS

the specification of which ☐ is attached hereto or ☒ was filed on December 6, 1990 as Application Serial No. 624,120 and was amended on _____ (if applicable).

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a). I claim foreign priority benefits under Title 35, United States Code, §112 or any foreign application(s) for patent or inventor's certificate listed below and have also identified _____ any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

COUNTRY	APPLICATION NUMBER	DATE OF FILING	PRIORITY CLAIMED UNDER 35 U.S.C. 119
			Yes _____ No _____
			Yes _____ No _____

I claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	DATE OF FILING	STATUS
<u>492,462</u>	<u>March 7, 1990</u>	<input type="checkbox"/> Patented <input checked="" type="checkbox"/> Pending <input type="checkbox"/> Abandoned
<u>362,901</u>	<u>June 7, 1989</u>	<input type="checkbox"/> Patented <input checked="" type="checkbox"/> Pending <input type="checkbox"/> Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) who are partners and associates in the firm of Townsend and Townsend to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Vern Norviel, Reg. No. 32,483

William M. Smith, Reg. No. 30,223

SEND CORRESPONDENCE TO: Vern Norviel

TOWNSEND and TOWNSEND

Steuart Street Tower, One Market Plaza
San Francisco, CA 94105

DIRECT TELEPHONE CALLS TO:

(name, registration number, and telephone number)

Vern Norviel, Reg. No. 32,483

☐ (415) 543-9600 or ☒ (415) 326-2400

201	FULL NAME OF INVENTOR	First Name	Middle Name or Initial
	RESIDENCE	City	State or Foreign Country
	CITIZENSHIP	City	Country of Citizenship
202	POST OFFICE ADDRESS	City	State or Country
	RESIDENCE	City	State or Country
	CITIZENSHIP	City	Country of Citizenship
203	POST OFFICE ADDRESS	City	State or Country
	RESIDENCE	City	State or Country
	CITIZENSHIP	City	Country of Citizenship

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 201	Signature of Inventor 202	Signature of Inventor 203
<i>Stephen P.A. Fodor</i>	<i>Lubert Stryer</i>	<i>James Winkler</i>
Date <u>JAN 28, 1991</u>	Date <u>Jan 31/1991</u>	Date <u>1/24/91</u>

DECLARATION AND POWER OF ATTORNEY.

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

IMPROVED POLYMER SYNTHESIS

The specification of which ☐ is attached hereto or ☒ was filed on December 6, 1990 as Application Serial No. 824,300 and was amended on _____ (if applicable)

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a). I claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

COUNTRY	APPLICATION NUMBER	DATE OF FILING	PRIORITY CLAIMED UNDER 35 U.S.C. 119
			Yes _____ No _____
			Yes _____ No _____

I claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	DATE OF FILING	STATUS
492,462	March 7, 1990	<input type="checkbox"/> Patented <input checked="" type="checkbox"/> Pending <input type="checkbox"/> Abandoned
362,901	June 7, 1989	<input type="checkbox"/> Patented <input checked="" type="checkbox"/> Pending <input type="checkbox"/> Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) who are partners and associates in the firm of Townsend and Townsend to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Vern Norviel, Reg. No. 32,483

William M. Smith, Reg. No. 30,223

SEND CORRESPONDENCE TO:		Vern Norviel TOWNSEND and TOWNSEND Steuart Street Tower, One Market Plaza San Francisco, CA 94105		DIRECT TELEPHONE CALLS TO: (name, registration number, and telephone number) Vern Norviel, Reg. No. 32,483 <input type="checkbox"/> (415) 543-9600 or <input checked="" type="checkbox"/> (415) 326-2400	
201	FULL NAME OF INVENTOR	Last Name HOLMES	First Name CHRISTOPHER	Middle Name or Initial P.	
	RESIDENCE & CITIZENSHIP	City Sunnyvale	State or Foreign Country California	Country of Citizenship U.S.A.	
	POST OFFICE ADDRESS	Post Office Address 521 Pine Avenue,	City Sunnyvale,	State or Country California	Zip Code 94086
202	FULL NAME OF INVENTOR	Last Name SOLAS	First Name DENNIS	Middle Name or Initial W.	
	RESIDENCE & CITIZENSHIP	City San Francisco	State or Foreign Country California	Country of Citizenship U.S.A.	
	POST OFFICE ADDRESS	Post Office Address 50 Gardenside Dr., #13,	City San Francisco,	State or Country California	Zip Code 94131
203	FULL NAME OF INVENTOR	Last Name	First Name	Middle Name or Initial	
	RESIDENCE & CITIZENSHIP	City	State or Foreign Country	Country of Citizenship	
	POST OFFICE ADDRESS	Post Office Address	City	State or Country	Zip Code

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 201 <i>Christopher P. Holmes</i>	Signature of Inventor 202 <i>Vern Norviel</i>	Signature of Inventor 203
Date <u>1/23/91</u>	Date <u>1/24/91</u>	Date

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Fodor, Stephen P.A.
Stryer, Lubert
Winkler, James L.
Holmes, Christopher P.
Solas, Dennis W.
- (ii) TITLE OF INVENTION: Very Large Scale Immobilized Polymer
Synthesis
- (iii) NUMBER OF SEQUENCES: 21
- (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Vernon A. Norviel
(B) STREET: One Market Plaza, Steuart Tower, Suite 2000
(C) CITY: San Francisco
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 94105
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE: 06-JUN-1995
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Norviel, Vernon A.
(B) REGISTRATION NUMBER: 32,483
(C) REFERENCE/DOCKET NUMBER: 16528J-000126
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 415-326-2400
(B) TELEFAX: 415-326-2422

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Tyr Gly Gly Phe Leu
1 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
220

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Pro Gly Gly Phe Leu
1 5

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Tyr Gly Ala Gly Phe
1 5

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Tyr Gly Ala Phe Leu Ser
1 5

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Tyr Gly Ala Phe Ser
1 5

[73]
[220]
[210]
[212]
[213]
[214]
[215]
[216]
[217]
[218]
[219]
[220]
[221]
[222]
[223]
[224]
[225]
[226]
[227]
[228]
[229]
[230]
[231]
[232]
[233]
[234]
[235]
[236]
[237]
[238]
[239]
[240]
[241]
[242]
[243]
[244]
[245]
[246]
[247]
[248]
[249]
[250]
[251]
[252]
[253]
[254]
[255]
[256]
[257]
[258]
[259]
[260]
[261]
[262]
[263]
[264]
[265]
[266]
[267]
[268]
[269]
[270]
[271]
[272]
[273]
[274]
[275]
[276]
[277]
[278]
[279]
[280]
[281]
[282]
[283]
[284]
[285]
[286]
[287]
[288]
[289]
[290]
[291]
[292]
[293]
[294]
[295]
[296]
[297]
[298]
[299]
[300]
[301]
[302]
[303]
[304]
[305]
[306]
[307]
[308]
[309]
[310]
[311]
[312]
[313]
[314]
[315]
[316]
[317]
[318]
[319]
[320]
[321]
[322]
[323]
[324]
[325]
[326]
[327]
[328]
[329]
[330]
[331]
[332]
[333]
[334]
[335]
[336]
[337]
[338]
[339]
[340]
[341]
[342]
[343]
[344]
[345]
[346]
[347]
[348]
[349]
[350]
[351]
[352]
[353]
[354]
[355]
[356]
[357]
[358]
[359]
[360]
[361]
[362]
[363]
[364]
[365]
[366]
[367]
[368]
[369]
[370]
[371]
[372]
[373]
[374]
[375]
[376]
[377]
[378]
[379]
[380]
[381]
[382]
[383]
[384]
[385]
[386]
[387]
[388]
[389]
[390]
[391]
[392]
[393]
[394]
[395]
[396]
[397]
[398]
[399]
[400]
[401]
[402]
[403]
[404]
[405]
[406]
[407]
[408]
[409]
[410]
[411]
[412]
[413]
[414]
[415]
[416]
[417]
[418]
[419]
[420]
[421]
[422]
[423]
[424]
[425]
[426]
[427]
[428]
[429]
[430]
[431]
[432]
[433]
[434]
[435]
[436]
[437]
[438]
[439]
[440]
[441]
[442]
[443]
[444]
[445]
[446]
[447]
[448]
[449]
[450]
[451]
[452]
[453]
[454]
[455]
[456]
[457]
[458]
[459]
[460]
[461]
[462]
[463]
[464]
[465]
[466]
[467]
[468]
[469]
[470]
[471]
[472]
[473]
[474]
[475]
[476]
[477]
[478]
[479]
[480]
[481]
[482]
[483]
[484]
[485]
[486]
[487]
[488]
[489]
[490]
[491]
[492]
[493]
[494]
[495]
[496]
[497]
[498]
[499]
[500]
[501]
[502]
[503]
[504]
[505]
[506]
[507]
[508]
[509]
[510]
[511]
[512]
[513]
[514]
[515]
[516]
[517]
[518]
[519]
[520]
[521]
[522]
[523]
[524]
[525]
[526]
[527]
[528]
[529]
[530]
[531]
[532]
[533]
[534]
[535]
[536]
[537]
[538]
[539]
[540]
[541]
[542]
[543]
[544]
[545]
[546]
[547]
[548]
[549]
[550]
[551]
[552]
[553]
[554]
[555]
[556]
[557]
[558]
[559]
[560]
[561]
[562]
[563]
[564]
[565]
[566]
[567]
[568]
[569]
[570]
[571]
[572]
[573]
[574]
[575]
[576]
[577]
[578]
[579]
[580]
[581]
[582]
[583]
[584]
[585]
[586]
[587]
[588]
[589]
[590]
[591]
[592]
[593]
[594]
[595]
[596]
[597]
[598]
[599]
[600]
[601]
[602]
[603]
[604]
[605]
[606]
[607]
[608]
[609]
[610]
[611]
[612]
[613]
[614]
[615]
[616]
[617]
[618]
[619]
[620]
[621]
[622]
[623]
[624]
[625]
[626]
[627]
[628]
[629]
[630]
[631]
[632]
[633]
[634]
[635]
[636]
[637]
[638]
[639]
[640]
[641]
[642]
[643]
[644]
[645]
[646]
[647]
[648]
[649]
[650]
[651]
[652]
[653]
[654]
[655]
[656]
[657]
[658]
[659]
[660]
[661]
[662]
[663]
[664]
[665]
[666]
[667]
[668]
[669]
[670]
[671]
[672]
[673]
[674]
[675]
[676]
[677]
[678]
[679]
[680]
[681]
[682]
[683]
[684]
[685]
[686]
[687]
[688]
[689]
[690]
[691]
[692]
[693]
[694]
[695]
[696]
[697]
[698]
[699]
[700]
[701]
[702]
[703]
[704]
[705]
[706]
[707]
[708]
[709]
[710]
[711]
[712]
[713]
[714]
[715]
[716]
[717]
[718]
[719]
[720]
[721]
[722]
[723]
[724]
[725]
[726]
[727]
[728]
[729]
[730]
[731]
[732]
[733]
[734]
[735]
[736]
[737]
[738]
[739]
[740]
[741]
[742]
[743]
[744]
[745]
[746]
[747]
[748]
[749]
[750]
[751]
[752]
[753]
[754]
[755]
[756]
[757]
[758]
[759]
[760]
[761]
[762]
[763]
[764]
[765]
[766]
[767]
[768]
[769]
[770]
[771]
[772]
[773]
[774]
[775]
[776]
[777]
[778]
[779]
[780]
[781]
[782]
[783]
[784]
[785]
[786]
[787]
[788]
[789]
[790]
[791]
[792]
[793]
[794]
[795]
[796]
[797]
[798]
[799]
[800]
[801]
[802]
[803]
[804]
[805]
[806]
[807]
[808]
[809]
[810]
[811]
[812]
[813]
[814]
[815]
[816]
[817]
[818]
[819]
[820]
[821]
[822]
[823]
[824]
[825]
[826]
[827]
[828]
[829]
[830]
[831]
[832]
[833]
[834]
[835]
[836]
[837]
[838]
[839]
[840]
[841]
[842]
[843]
[844]
[845]
[846]
[847]
[848]
[849]
[850]
[851]
[852]
[853]
[854]
[855]
[856]
[857]
[858]
[859]
[860]
[861]
[862]
[863]
[864]
[865]
[866]
[867]
[868]
[869]
[870]
[871]
[872]
[873]
[874]
[875]
[876]
[877]
[878]
[879]
[880]
[881]
[882]
[883]
[884]
[885]
[886]
[887]
[888]
[889]
[890]
[891]
[892]
[893]
[894]
[895]
[896]
[897]
[898]
[899]
[900]
[901]
[902]
[903]
[904]
[905]
[906]
[907]
[908]
[909]
[910]
[911]
[912]
[913]
[914]
[915]
[916]
[917]
[918]
[919]
[920]
[921]
[922]
[923]
[924]
[925]
[926]
[927]
[928]
[929]
[930]
[931]
[932]
[933]
[934]
[935]
[936]
[937]
[938]
[939]
[940]
[941]
[942]
[943]
[944]
[945]
[946]
[947]
[948]
[949]
[950]
[951]
[952]
[953]
[954]
[955]
[956]
[957]
[958]
[959]
[960]
[961]
[962]
[963]
[964]
[965]
[966]
[967]
[968]
[969]
[970]
[971]
[972]
[973]
[974]
[975]
[976]
[977]
[978]
[979]
[980]
[981]
[982]
[983]
[984]
[985]
[986]
[987]
[988]
[989]
[990]
[991]
[992]
[993]
[994]
[995]
[996]
[997]
[998]
[999]
[1000]

[illegible]

(A) LENGTH: 5 amino acids

(C) STRANDEDNESS: single

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Tyr Gly Ala Phe Leu

1

5

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Tyr Gly Gly Phe Leu Ser
1 5

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Tyr Gly Ala Phe

1

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Tyr Gly Ala Leu Ser
1 5

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Tyr Gly Gly Phe Ser
1 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
220

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Tyr Gly Ala Leu

1

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Tyr Gly Ala Phe Leu Phe
1 5

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Tyr Gly Ala Phe Phe
1 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Tyr Gly Gly Leu Ser
1 5

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Tyr Gly Gly Phe Leu
1 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
22

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Tyr Gly Ala Phe Ser Phe
1 5

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Tyr Gly Ala Phe Met Gln
1 5

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Tyr Gly Ala Phe Met
1 5

[1]
[2]
[3]
[4]
[5]
[6]
[7]
[8]
[9]
[10]
[11]
[12]
[13]
[14]
[15]
[16]
[17]
[18]
[19]
[20]
[21]
[22]
[23]
[24]
[25]
[26]
[27]
[28]
[29]
[30]
[31]
[32]
[33]
[34]
[35]
[36]
[37]
[38]
[39]
[40]
[41]
[42]
[43]
[44]
[45]
[46]
[47]
[48]
[49]
[50]
[51]
[52]
[53]
[54]
[55]
[56]
[57]
[58]
[59]
[60]
[61]
[62]
[63]
[64]
[65]
[66]
[67]
[68]
[69]
[70]
[71]
[72]
[73]
[74]
[75]
[76]
[77]
[78]
[79]
[80]
[81]
[82]
[83]
[84]
[85]
[86]
[87]
[88]
[89]
[90]
[91]
[92]
[93]
[94]
[95]
[96]
[97]
[98]
[99]
[100]

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Tyr Gly Ala Phe Gln
1 5

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Tyr Gly Gly Phe Met
1 5